

Sodium toxicity (proportion of land affected)

High sodicity is one of the characteristics linked to the strong marine influence on many southern South Australian soils


Sodium toxicity occurs mostly in the drier parts of Southern South Australia where there is a very high level of deep subsoil sodicity (i.e. exchangeable sodium percentages (ESP) exceeding 25). This generally occurs at depths of between 50 and 100 cm, sometimes shallower. These conditions are invariably associated with high pH, moderate salinity and often high boron concentrations, all of which are natural features of these soils. They are not necessarily associated with poor soil structure. There is some evidence to suggest that these high levels of sodicity are toxic to some plants, particularly horticultural species. If there are toxic effects, it is reasonable to assume that in some years, the sodicity is preventing optimum water use efficiency, and could therefore be contributing to rising watertables. In practice, neither gypsum nor any other ameliorant will have any significant effect on these soils, due to the very high rates required and the difficulty of incorporation to the required depth. The use of plants with sodium tolerance is one solution. This implies cultivar selection focused on sodium tolerance or the use of naturally occurring plants with sodium tolerance.

Land assessment in southern South Australia

Estimates of ESP are based on extrapolation of laboratory analyses between similar soil materials and soil types.

Soil properties can vary across the landscape in a subtle or dramatic fashion. [Mapping at a regional scale](#) is not able to display this level of variability, however proportions of each *Sodium toxicity (proportion of land affected)* class (e.g. TNAP1, TNAP2, etc.) have been estimated for each map unit.

Further information can be found in [Assessing Agricultural Land](#) (Maschmedt 2002).



Poorly structured sodic subsoil from 15 cm over sodium toxic clay (exchangeable sodium percentage of 51) from 35 cm

Area statistics

Area affected by sodium toxicity (exchangeable sodium percentages >25 shallower than 100 cm)	Area	Cleared land	Class*
Land not affected by sodium toxicity	56.06%	52.28%	TNAP1
Land affected by sodium toxicity	42.52%	46.01%	TNAP2
Not applicable	1.43%	1.70%	TNAPX
TOTAL HECTARES	15,765,460	10,439,300	

* The letters 'TNAP' denotes classes that are specific to *Sodium toxicity (proportion of land affected)*




Displaying data in soil maps

Soil and land attribute maps display a simplified version of the underlying data. Mapping classes are based on an interpretation of soil landscape map units within which the depth to highly sodic layers can vary. Each map unit component is assessed according to the estimated average depth to toxic sodium (ESP exceeding 25). Map units are then classified according to the proportion of the map unit area where this occurs within the top 100 cm of soil.



Saline land with near-surface sodium toxicity



Further information

- View data on [NatureMaps](#) (→ Soils)
- Read the [metadata](#) for this layer
- Read more about [soil attribute mapping](#)
- Contact [Mapland](#)

Download from Enviro Data SA:

- [Statewide map](#) and [spatial dataset](#)
- [Assessing Agricultural Lands](#) (Maschmedt 2002)
- Soils of Southern SA book [Part 1](#) and [Part 2](#)



This work is licensed under the Creative Commons Attribution 4.0 International License.

To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© Crown in right of the State of South Australia, through the Department of Environment, Water and Natural Resources 2016. Last updated June 2016.

