

Great Australian Bight

Fowlers Bay

Ceduna

Streaky Bay

Kimba

Port Augusta

Whyalla

Port Pirie

Hawker

Quorn

Jamestown

Elliston

Lock

Cleve

Kadina

Moonta

Burra

Clare

Cummins

Port Lincoln

Spencer Gulf

Maitland

Morgan

Nuriootpa

Renmark

Berri

Loxton

Yorke town

ADELAIDE

Gawler

Mt Barker

Murray Bridge

Strathalbyn

Pinnaroo

Gulf St Vincent

Victor Harbor

Meningie

Keith

Kingscote

Kingston

Robe

Naracoorte

Penola

Mt Gambier

NEW SOUTH WALES

VICTORIA

SOUTHERN SOUTH AUSTRALIA

Land Use Potential for CHESTNUTS

Potential based on soil and landscape attributes only - no account has been taken of water quality or availability, climatic factors or existing land use

Proportion of land with moderate to high potential

Most common potential class

	More than 60%	High
	More than 60%	Moderately high
	More than 60%	Moderate to high
	More than 60%	Moderate
	30-60%	Low to high
	10-30%	Moderately low to low
	1-10%	Moderately low to low
	Less than 1%	Moderately low
	Less than 1%	Low
	Not applicable	

PRELIMINARY MAP

HIGH POTENTIAL: Land with high productive potential and requiring no more than standard management practices to sustain productivity.

MODERATELY HIGH POTENTIAL: Land with moderately high productive potential and / or requiring specific, but widely accepted and used management practices to sustain productivity.

MODERATE POTENTIAL: Land with moderate productive potential and / or requiring specialised management practices to sustain productivity.

MODERATELY LOW POTENTIAL: Land with marginal productive potential and / or requiring very highly specialised management skills to sustain productivity.

LOW POTENTIAL: Land with low productive potential and / or permanent limitations which effectively preclude its use.

NOTES ON USE OF THE MAP:

1. Classes are based on interpretations of Soil Landscape Map Units. The most limiting feature of a Soil Landscape Map Unit determines the overall class of that unit. Soil Landscape Map Units are not homogenous entities – the class is intended to reflect the most common characteristics of the landscape. Unspecified variations occur.
2. Boundaries between mapping units should be treated as transition zones.
3. This information is derived from limited field assessments and / or laboratory data, and estimates based on personal experience or judgement may be used where data are unavailable.
4. The interpretation methodologies are in developmental stage, and only limited verification has been undertaken. Mapping classes are subject to change without notice.
5. The map is intended to provide a regional overview and should not be used to draw conclusions about specific locations.
6. The scale of maps should not be enlarged beyond their scale of publication.
7. Independent expert advice should be sought prior to using this information for commercial decision-making.



Licensed under Creative Commons Attribution 4.0 International License
<http://creativecommons.org/licenses/by/4.0/>

© Crown in right of the State of South Australia
Department for Environment and Water, 2018

Land assessment: DEW Soil and Land Program, 2009
Map projection: Lambert Conformal Conic
Map datum: GDA94



Government of South Australia

Department for Environment
and Water



0 50 100 150 200

Kilometres

— Main road

Southern Ocean