

Great Australian Bight

Fowlers Bay

Ceduna

Streaky Bay

Kimba

Whyalla

Hawker

Quorn

Port Augusta

Port Pirie

Jamestown

Burra

Clare

Morgan

Kadina
Moonta

Maitland

Nuriootpa

Gawler

Yorke town

ADELAIDE

Mt Barker

Murray Bridge

Strathalbyn

Victor Harbor

Meningie

Renmark

Berri

Loxton

Pinnaroo

Kingscote

Keith

Kingston

Naracoorte

Robe

Penola

Mt Gambier

NEW SOUTH WALES

VICTORIA

SOUTHERN SOUTH AUSTRALIA

ACID SULFATE SOIL POTENTIAL

Classes are based on an interpretation of soil landscape map units which are classified according to the proportion of area susceptible to the development of acid sulfate soils.

Proportion of land susceptible to the development of acid sulfate soils

- Negligible
- 1-10%
- 10-30%
- 30-60%
- More than 60%
- Incomplete data (usually wet inland areas)
- Not applicable

NOTES ON USE OF THE MAP

- The information is derived from limited field inspection, and is subject to change without notice.
- Boundaries between mapping units should be treated as transition zones.
- The map is intended to provide a regional overview and should not be used to draw conclusions about conditions at specific locations.
- The scale of maps should not be enlarged beyond their scale of publication.
- Independent expert advice should be sought prior to using this information for commercial decision-making.



Licensed under Creative Commons Attribution 4.0 International License
<http://creativecommons.org/licenses/by/4.0/>

© Crown in right of the State of South Australia
Department for Environment and Water, 2018

Land assessment: DEW Soil and Land Program, 2009
Map projection: Lambert Conformal Conic
Map datum: GDA94



0 50 100 150 200

Kilometres

Main road

Southern Ocean



Government of South Australia

Department for Environment
and Water