

Great Australian Bight

Fowlers Bay

Ceduna

Streaky Bay

Kimba

Whyalla

Hawker

Quorn

Port Augusta

Port Pirie

Jamestown

Elliston

Lock

Cleve

Burra

Kadina
Moonta

Clare

Morgan

Cummins

Maitland

Nuriootpa
Gawler

Renmark

Berri

Loxton

Port Lincoln

Spencer Gulf

Gulf St Vincent

Yorketown

ADELAIDE

Mt Barker

Murray Bridge

Strathalbyn

Pinnaroo

Kingscote

Victor Harbor

Meningie

Keith

Kingston

Robe

Mt Gambier

Penola

Naracoorte

NEW SOUTH WALES

VICTORIA

Southern Ocean

SOUTHERN SOUTH AUSTRALIA

INHERENT FERTILITY

Classes are based on an interpretation of soil landscape map units which are classified according to the inherent fertility of the component soils, on a weighted average basis. Inherent fertility rankings are assessed according to soil texture, leaching capacity, acidification potential, carbonate and ironstone content.

- High to very high
- Moderate
- Moderately low
- Low
- Very low
- Not applicable

NOTES ON USE OF THE MAP

1. The information is derived from limited field inspection, and is subject to change without notice.
2. Boundaries between mapping units should be treated as transition zones.
3. The map is intended to provide a regional overview and should not be used to draw conclusions about conditions at specific locations.
4. The scale of maps should not be enlarged beyond their scale of publication.
5. Independent expert advice should be sought prior to using this information for commercial decision-making.



Licensed under Creative Commons Attribution 4.0 International License
<http://creativecommons.org/licenses/by/4.0/>

© Crown in right of the State of South Australia
 Department for Environment and Water, 2018

Land assessment: DEW Soil and Land Program, 2009
 Map projection: Lambert Conformal Conic
 Map datum: GDA84



0 50 100 150 200

Kilometres

— Main road



Government of South Australia

Department for Environment
and Water