
Great Australian Bight
Fowlers Bay
Ceduna
Streaky Bay
Kimba
Lock
Elliston
Cleve
Cummins
Port Lincoln
Whyalla
Hawker
Quorn
Port Augusta
Port Pirie
Jamestown
Burra
Clare
Kadina
Moonta
Morgan
Maitland
Spencer Gulf
Nuriootpa
Gawler
Renmark
Berri
Loxton
ADELAIDE
Mt Barker
Gulf St Vincent
Yorketown
Murray Bridge
Strathalbyn
Pinnaroo
Victor Harbor
Kingscote
Meningie
Keith
Kingston
Naracoorte
Robe
Penola
Mt Gambier
Southern Ocean
NEW SOUTH WALES
VICTORIA
SOUTHERN SOUTH AUSTRALIA
ROOTZONE DEPTH POTENTIAL
Horticultural crop type CC -
Hardy crops (eg grapes, olives)
Classes are based on an interpretation of soil landscape map units which are classified according to the estimated potential rootzone depth of the component soils, on a weighted average basis.
Average potential rootzone depth
More than 100 cm
80-100 cm
60-80 cm
50-60 cm
40-50 cm
30-40 cm
20-30 cm
Less than 20 cm
Not applicable
NOTES ON USE OF THE MAP
1. The information is derived from limited field inspection, and is subject to change without notice.
2. Boundaries between mapping units should be treated as transition zones.
3. The map is intended to provide a regional overview and should not be used to draw conclusions about conditions at specific locations.
4. The scale of maps should not be enlarged beyond their scale of publication.
5. Independent expert advice should be sought prior to using this information for commercial decision-making.
N
0 50 100 150 200
Kilometres
Main road
Licensed under Creative Commons Attribution 4.0 International License
http://creativecommons.org/licenses/by/4.0/
© Crown in right of the State of South Australia
Department for Environment and Water, 2018
Land assessment: DEW Soil and Land Program, 2009
Map projection: Lambert Conformal Conic
Map datum: GDA94
Government of South Australia
Department for Environment and Water