

Great Australian Bight

Fowlers Bay

Ceduna

Streaky Bay

Kimba

Whyalla

Hawker

Quorn

Port Augusta

Port Pirie

Jamestown

Elliston

Lock

Cleve

Kadina  
Moonta

Clare

Burra

Cummins

Port Lincoln

Maitland

Morgan

Nuriootpa

Gawler

Yorketown

Gulf St Vincent

Adelaide

Murray Bridge

Strathalbyn

Victor Harbor

Meningie

Renmark

Berri

Loxton

Kingscote

Keith

Kingston

Naracoorte

Robe

Penola

Mt Gambier

NEW SOUTH WALES


VICTORIA

## SOUTHERN SOUTH AUSTRALIA

### SALINITY - Non-watertable (magnesia patches)

Classes are based on an interpretation of soil landscape map units. Map units are classified according to the proportion of area affected by magnesia patches.

Proportion of land affected by magnesia patches



#### NOTES ON USE OF THE MAP


1. The information is derived from limited field inspection, and is subject to change without notice.
2. Boundaries between mapping units should be treated as transition zones.
3. The map is intended to provide a regional overview and should not be used to draw conclusions about conditions at specific locations.
4. The scale of maps should not be enlarged beyond their scale of publication.
5. Independent expert advice should be sought prior to using this information for commercial decision-making.



Licensed under Creative Commons Attribution 4.0 International License  
<http://creativecommons.org/licenses/by/4.0/>

© Crown in right of the State of South Australia  
Department for Environment and Water, 2018

Land assessment: DEW Soil and Land Program, 2009  
Map projection: Lambert Conformal Conic  
Map datum: GDA84



Main road

Southern Ocean



Government of South Australia

Department for Environment  
and Water