

Great Australian Bight

Fowlers Bay

Ceduna

Streaky Bay

Kimba

Whyalla

Hawker

Quorn

Port Augusta

Port Pirie

Jamestown

Burra

Clare

Kadina
Moonta

Maitland

Nuriootpa
Gawler

Morgan

Renmark

Berri

Loxton

Spencer Gulf

Port Lincoln

Gulf St Vincent

Yorketown

ADELAIDE

Mt Barker

Murray Bridge

Strathalbyn

Victor Harbor

Meningie

Keith

Naracoorte

Kingston

Robe

Penola

Mt Gambier

NEW SOUTH WALES

VICTORIA

SOUTHERN SOUTH AUSTRALIA

SODIUM TOXICITY (proportion of land affected)

Classes are based on an interpretation of soil landscape map units which are classified according to the proportion of area with exchangeable sodium percentages (ESP) exceeding 25 in the upper 100 cm of the soil.

Proportion of land with sodium toxicity in the upper 100 cm of soil.

- Negligible
- 1-10%
- 10-30%
- 30-60%
- More than 60%
- Not applicable

NOTES ON USE OF THE MAP

1. The information is derived from limited field inspection, and is subject to change without notice.
2. Boundaries between mapping units should be treated as transition zones.
3. The map is intended to provide a regional overview and should not be used to draw conclusions about conditions at specific locations.
4. The scale of maps should not be enlarged beyond their scale of publication.
5. Independent expert advice should be sought prior to using this information for commercial decision-making.



Licensed under Creative Commons Attribution 4.0 International License
<http://creativecommons.org/licenses/by/4.0/>

© Crown in right of the State of South Australia
Department for Environment and Water, 2018

Land assessment: DEW Soil and Land Program, 2009
Map projection: Lambert Conformal Conic
Map datum: GDA84



0 50 100 150 200

Kilometres

— Main road

Southern Ocean



Government of South Australia

Department for Environment
and Water