

Benthic Macroinvertebrate Response Monitoring in the Coorong and Murray Mouth, 2013/14



Final Report **for the Department of Environment and Natural Resources**

Sabine Dittmann, Tanith Ramsdale, Nat Navong and Angela McGuire

Flinders University, School of Biological Sciences



Flinders
UNIVERSITY



Government of South Australia
Department of Environment
and Natural Resources

This report may be cited as: Dittmann, S., Ramsdale, T., Navong, N. and McGuire, A., 2014: Benthic Macroinvertebrate Response Monitoring in the Coorong and Murray Mouth, 2013/14. Report for the Department of Environment, Water and Natural Resources, Adelaide.

Contents

1. Executive Summary	1
2. Introduction	5
3. Materials and Methods	7
3.1 Sampling sites and dates	7
3.2 Environmental parameters	9
3.3 Macrofauna	10
3.4 Data analysis	10
4. Results and Discussion	12
4.1 Environmental Conditions	12
4.2 Macroinvertebrate Recruitment	15
4.3 Macroinvertebrate Recolonisation	24
4.4 Relationships between Macroinvertebrate Communities and Environmental Conditions	40
5 Conclusions	46
6 Acknowledgements	47
7 References	47

1. Executive Summary

- This report presents findings from monitoring the response of macroinvertebrates in the Murray Mouth and Coorong in the fourth year since flow resumed in late 2010. All methods used followed procedures as during previous monitoring. The nine mudflat sites were located between Monument Road near the Goolwa barrage, and Villa de Yumpa at the northern end of the South Lagoon. Sampling occurred on three occasions between late November 2013 and March 2014, with the sampling in December 2013 also covering the subtidal sediments and two locations on the Younghusband Peninsula.
- The monitoring centred on a series of key questions on indications for continued recovery of this estuarine and lagoon system (Table 1). Following indications of a recovery in the previous monitoring period, the predictions for this current survey had been that further recolonisation would occur and populations of macroinvertebrates re-establish in mudflat of the Murray Mouth and Coorong.
- To establish recovery of macroinvertebrate populations and communities, measurements were taken to assess species richness, abundance and size-frequencies. To explore for evidence of how macroinvertebrate communities are affected by environmental conditions, water and sediment variables were determined on each sampling occasion.
- Signs of recovery were based on higher numbers of species and increased abundances of species like *Simplisetia aequisetis* at multiple sites throughout the Murray Mouth and in the northern North Lagoon. Most species maintained their distribution ranges as described in the 2012/13 monitoring, and single individuals of more species were found around Hells Gate and the northern South Lagoon. Species like the small bivalve *Arthritica helmsi*, which was very abundant before the drought, continued to recolonise the mudflats. Small hydrobiid snails became abundant as well and larger bivalves (*Soletellina alba*, *Spisula trigonella*) were found at several sites. The findings from the monitoring indicate an increasing complexity of the macroinvertebrate community with more functional richness, such as increased bioturbation in sediments.
- Macroinvertebrate abundances were mostly higher in subtidal sediments than intertidal mudflats during the November/December 2013 survey. In February and March 2014, abundances in the mudflats were substantially reduced, reflecting a decrease in numbers of amphipods and chironomid larvae. As subtidal sediments were not sampled at these further surveys, it is unknown whether abundances remained high in the subtidal or not. The lack of further sampling in the subtidal also affected the assessment of population sizes for several species.
- Size frequency distributions gave indications of recruitment for most species of macroinvertebrates and signs of continuous recruitment over the summer period for some polychaete species. Juvenile *S. aequisetis* and *N. australiensis* were more frequent in sub- than intertidal sediments in December 2013.
- The macroinvertebrate communities in the Murray Mouth and Coorong during the 2013/14 monitoring were similar to the previous post-flow period. In the Murray Mouth, several analyses indicated a return to pre-drought conditions, while the trajectory for the Coorong may lead into a new direction. Clear community differences for pre-drought, drought and flow periods were identified. In the current monitoring, variations in macroinvertebrate communities were explained mainly by salinity and sediment compositions.
- This monitoring documented that macroinvertebrate populations were re-establishing in the Murray Mouth and Coorong, yet the unexplained decrease in abundances over summer 2013/14 could have affected further recovery. More comprehensive sampling of inter- and subtidal sediments in

the future may ascertain whether the subtidal sediments can function as a source for mudflat recolonisation and thus support the resilience of this ecosystem.

Table 1. Summary table of key questions and findings for the 2013/2014 macroinvertebrate surveys in response to continued flow.

Key Questions	Summary of Findings
<p>1. Are there indications of continued system recovery in 2013/14 following the significant flows of 2010-11 and further flows in 2011, 2012 and 2013?</p>	<ul style="list-style-type: none"> • Yes, monitoring revealed improved estuarine conditions in the Murray Mouth and North Lagoon and macroinvertebrate communities started to re-establish. This was particularly pronounced in late spring/early summer, while abundances of several benthic taxa decreased over the course of summer for unknown reasons.
<p>a) Will environmental conditions within the Murray Mouth reflect true estuarine conditions and have they further improved in the North and South Lagoons since flows recommenced in 2010?</p>	<ul style="list-style-type: none"> • Seasonal changes in salinity occurred in response to the volume of flow across the barrages, with lower salinities recorded in late spring/early summer, and increasing salinities over summer when flow volumes decreased. Seasonal variations in salinity subject to river flow are typical for estuaries, and estuarine conditions were thus improved in the Murray Mouth and North Lagoon. Salinities remained hypersaline in the South Lagoon. • While the salinity gradient from brackish to marine conditions in the Murray Mouth to hypersaline conditions in the South Lagoon persisted, salinities were >30 ppt lower in each region than during the drought. • In the Murray Mouth, salinities changed from freshwater during the 2010/11 flow, to brackish and marine in the following monitoring periods. Salinities in the 2013/14 monitoring period were comparable to the previous 2012/13 monitoring. • Salinities were slightly higher at the subtidal and peninsula locations during sampling in December, yet no vertical stratification of salinity was recorded.
<p>b) Has recruitment continued for macroinvertebrate species that began recovery in 2011-12 and is recruitment occurring for other species in 2013-14?</p>	<ul style="list-style-type: none"> • Recruitment continued for most macroinvertebrate species as in the previous survey, with additional recruitment of <i>Arthritica helmsi</i> recorded in this survey. • Small individuals were encountered for most macroinvertebrate species in the Murray Mouth and North Lagoon, and amphipods carrying eggs were seen in the December samples. • Recruitment continued in summer 2013/14 for the polychaetes <i>Simplisetia aequisetis</i>, <i>Boccardiella limnicola</i> and <i>Capitella</i> spp.. Other polychaetes (<i>Nephtys australiensis</i> and <i>Australonereis ehlersi</i>) occurred in too low numbers to evaluate recruitment, although small sized <i>N. australiensis</i> were found in December. • Juveniles of <i>S. aequisetis</i>, <i>Capitella</i> spp. and the small bivalve <i>A. helmsi</i> were present at all three sampling events, indicating continuous reproduction over summer. Size-frequency distributions for <i>B. limnicola</i> indicated a recruitment event in December. • The bivalve <i>Arthritica helmsi</i> was recorded with a similar size range over all three sampling events in the Murray Mouth and North Lagoon. Other unidentified juvenile bivalves occurred in December 2013 in the Murray Mouth. Small individuals of

Key Questions	Summary of Findings
	<p><i>Soletellina alba</i> were found in the Murray Mouth.</p> <ul style="list-style-type: none"> The occurrence of juveniles at sampling sites overlapped with the distribution of larger specimens. The frequency of small sized <i>S. aequisetis</i> and <i>N. australiensis</i> was higher in sub- than intertidal sediments.
<p>c) Can the recruitment of species in 2013/14 be linked with the timing of differing flow scenarios to identify drivers of macroinvertebrate recruitment?</p>	<ul style="list-style-type: none"> The presence of small-sized specimens of several macroinvertebrate species, in particular in December and February, gave an indication of recruitment occurring in early summer, as flow volumes decreased and salinities increased. The continuous recruitment of some macroinvertebrates indicates either that recruitment is triggered by other cues than those related to flow and salinity changes, or a life history adapted to the variable environment of estuaries.
<p>d) Will there be any further recolonisation of mudflats by macroinvertebrates absent from areas in previous monitoring 2010-12?</p>	<ul style="list-style-type: none"> The number of macroinvertebrate species found in mudflats was higher than in the previous survey. Macroinvertebrates that had previously colonised mudflats from subtidal locations had strengthened their population size in mudflats in the Murray Mouth and North Lagoon. The small bivalve, <i>Arthritica helmsi</i> continues to recolonise mudflat areas, with abundances increasing at several mudflat sites during the 2013/14 survey. Apart from Hunters Creek and Ewe Island, macroinvertebrate abundances were higher in the subtidal than intertidal mudflats in December 2013. Abundances in mudflats decreased substantially over summer 2013/14.
<p>e) Will recolonisation of macroinvertebrate species occur in the South Lagoon if salinities remain lowered in this region?</p>	<ul style="list-style-type: none"> Insect larvae (mainly chironomids and Ephydriidae) were still prominent in sediments at Parnka Point and Villa de Yumpa. Yet numerically, ostracods were more abundant at Parnka Point. Single individuals of estuarine macroinvertebrates were found, such as amphipods in December 2013 and March 2014, oligochaetes and the polychaete <i>Capitella spp.</i> in subtidal sediments at Villa de Yumpa, and <i>Arthritica helmsi</i> in the mudflat at this site in December 2013. Finding specimens of estuarine macroinvertebrates around Hells Gate may indicate that recolonisation may occur if environmental conditions continue to improve.
<p>f) Have species maintained or further increased their distribution range in comparison to 2011-12?</p>	<ul style="list-style-type: none"> Most species maintained their distribution range in comparison to previous expansions observed between 2012 and 2013. While species occurred again at the same sites, shifts in abundance across sites were recorded. There were only few indications for increased distribution ranges in 2013/14, such as <i>Capitella sp.</i> expanding into the Murray Mouth.
<p>g) Has the macroinvertebrate community been restored to pre-drought conditions following the barrage flows into the Coorong since late 2010 and are they</p>	<ul style="list-style-type: none"> In the Murray Mouth, a return to a macroinvertebrate community more similar to the pre-drought was indicated by the trajectory of community change over the last decade. In the Coorong, site and regional differences in community developments obscured a clear trend, yet change away from macroinvertebrate communities present during the drought was

Key Questions	Summary of Findings
comparable to values observed prior to drought conditions?	<p>apparent.</p> <ul style="list-style-type: none"> • The macroinvertebrate species characterising the communities in the Murray Mouth and Coorong were respectively corresponding communities for particular salinity regimes three decades ago.
2. Which environmental conditions influence the distribution, abundance and community structure of macroinvertebrates?	<ul style="list-style-type: none"> • Salinity was again found to be the key driver for the distribution, abundance and community structure of macroinvertebrates, with sediment factors (grain-size fractions, chlorophyll-a content) being secondary in influence over temporal and spatial changes in macroinvertebrate communities. • The relevance of salinity and sediment properties in explaining variation in macroinvertebrate communities was more pronounced in the Murray Mouth and South Lagoon than in the North Lagoon.
3. Are there similarities or differences in the community structure of macroinvertebrates across differing flow scenarios (drought/flood)?	<ul style="list-style-type: none"> • Dissimilarities were highest between the drought period in comparison to the pre-drought and flow periods. While some macroinvertebrate species were present in all periods, other were lacking during the drought or during the flow, or changes in their abundances affected their contribution to characterising the communities at the time. • The macroinvertebrate community in the current monitoring year 2013/14 was classified as similar to the other years of the flow period for both the Murray Mouth and the Coorong. • In the Murray Mouth, there were clear distinctions between macroinvertebrate community structure observed under drought and flow conditions, • In the Coorong, recovery was slower, but communities during periods of drought and flow were distinct to each other.

2. Introduction

Restored freshwater flows into the Murray Mouth and Coorong were predicted to lead to a recovery of this ecosystem, composed of estuarine and lagoon habitats. The flow that resumed in late 2010 provided the first freshwater input after a prolonged drought (Leblanc et al. 2012) and restored connectivity between the river, lake, estuary, lagoon and adjacent sea. Restoring the physico-chemical setting and boundary conditions has supported ecological recovery in coastal and estuarine ecosystems elsewhere (Elliott et al. 2007, Borja et al. 2010). Recovery can be idiosyncratic, affected by species traits and connectivity, and the types of pressures that occurred (see review by (Duarte et al. 2014)). In many examples worldwide, recovery is slow and only partial as thresholds differ for degradation and recovery phases and global change causes shifting baselines (Duarte et al. 2014). Prior disturbance history can also affect resilience of estuarine populations (Dolbeth et al. 2007). With recovery ecology receiving more recent scientific attention (Lotze et al. 2006, Verdonshot et al. 2013), monitoring the response of macroinvertebrates in the Murray Mouth and Coorong can not only advise management on the efficiency of flow restoration measures (Verissimo et al. 2012, Marques et al. 2013), but also contribute to the general understanding of processes for the recovery of benthic communities, which are an essential part of coastal and marine ecosystems (Levin et al. 2001).

This report presents findings from the fourth year of monitoring the response of macroinvertebrates to the restored freshwater inflows into the Murray Mouth and Coorong since late 2010 (Figure 1). While the volume of flow decreased substantially since the main flow event in 2010/11, a seasonal pattern of mainly winter/early spring flow over the barrages into the Murray Mouth and Coorong remained since 2011/12. The previous survey years gave indications of a recovery, following a slow start in the first year of inflow (Dittmann et al. 2013a). Species not seen in the system during the drought were first found again in the 2012/13 survey (Dittmann et al. 2013a), and their establishment will affect whether benthic communities return to pre-drought conditions, or follow a trajectory into a new state.

In this monitoring, we aim to evaluate whether the changed environmental conditions, expected to return to estuarine characteristics, are facilitating recovery of populations of macroinvertebrates, an increase in the number of species inhabiting the sediments, and a spread in the occurrence of species further into the southern reaches of the Coorong. The approach and methods followed previous monitoring protocols, but the design was changed with a reduction to three sampling events over summer 2013/14 and only one sampling event including subtidal (permanently submerged) sediments and intertidal (episodically exposed) mudflats on the peninsula side of the Coorong and Murray Mouth. Diversity, abundance and distribution ranges of macroinvertebrates were expected to be higher and larger in the more benign subtidal locations than the mudflats, where variability in exposure and temperature can affect small sized macroinvertebrates (Hummel et al. 1986, 1988).

The report is structured around a series of key questions (Table 1) and detailed data and outcome of statistical analyses provided in an appendix. The findings are presented as results and discussion, with a summary of key outcomes captured in Table 1. Some final conclusions and recommendations for further monitoring conclude the report.




Figure 1: Monthly barrage flow from the Lower Lakes into the Murray Mouth and Coorong from August 2010 up until April 2014, based on data from the MDBA. The asterisks indicate when sampling took place for monitoring the response of macroinvertebrates to the recommenced flow.

3. Materials and Methods

3.1 Sampling sites and dates

Three sampling events took place for the monitoring of water release effects and recovery in the Murray Mouth and Coorong in the fourth year since flow resumed. Sampling occurred in November/December 2013 (in conjunction with The Living Murray (TLM) condition monitoring for macroinvertebrates and mudflats, Dittmann et al. 2014), and in February and March 2014. Water quality was measured in the field and samples were taken for sediment characteristics, macrobenthic communities and size frequency distributions.

The ten sampling sites were located in the Murray Mouth (Monument Road (MR), Hunters Creek (HC), Ewe Island (EI), and Pelican Point (PP), North Lagoon of the Coorong (Mark Point (MP), Long Point (LP), Noonameena (NM)), and at Parnka Point (on the northern and southern side of Hells Gate, referred to as PaPN and PaPS)) and at Villa de Yumpa (VdY) in the South Lagoon (Figure 2). These sites were sampled in the previous studies for monitoring the effects of the water release, although not all sites were sampled every year in the past (Dittmann et al. 2011, Keuning et al. 2012, Dittmann et al. 2013a). Some of the sampling sites overlap with sites for TLM condition monitoring (see Dittmann et al. 2013b), allowing comparisons over longer time frames.




Figure 2. Study sites in the Murray Mouth and Coorong during the summer 2013/2014 monitoring period, see Table 2 for site acronyms. Sampling locations per site are indicated as open circles for ‘intertidal’, closed circles for submerged locations, and grey circles for peninsula locations (at Pelican Point and Long Point only). Parnka Point was sampled on the northern and southern side of Hells Gate on some occasions.

Table 2. Overview of the macroinvertebrate sampling in the Murray Mouth and Coorong in summer 2013/14, showing the dates and months of various sampling activities. Subtidal and peninsula sites were only sampled in December 2013. See Figure 2 for site locations.

Site	Sampling date		
	Nov/Dec 2013	Feb-14	Mar-14
Monument Road (MR) - Intertidal	25/11/2013	17/02/2014	17/03/2014
Subtidal	2/12/2013		
Hunters Creek (HC) - Intertidal	25/11/2013	17/02/2014	17/03/2014
Subtidal	2/12/2013		
Ewe Island (EI) - Intertidal	25/11/2013	17/02/2014	17/03/2014
Subtidal	2/12/2013		
Pelican Point (PP) - Intertidal	10/12/2013	17/02/2014	17/03/2014
Subtidal	3/12/2013		
Peninsula	3/12/2013		
Mark Point (MP) - Intertidal	10/12/2013	18/02/2014	19/03/2014
Subtidal	3/12/2013		
Long Point (LP) - Intertidal	10/12/2013	18/02/2014	19/03/2014
Subtidal	3/12/2013		
Peninsula	3/12/2013		
Noonameena (NM) - Intertidal	10/12/2013	18/02/2014	19/03/2014
Subtidal	3/12/2013		
Parnka Point North (PaPN) - Intertidal	9/12/2013		
Subtidal	3/12/2013		
Parnka Point South (PaPS) - Intertidal	9/12/2013	19/02/2014	19/03/2014
Villa de Yumpa (VdY) - Intertidal	9/12/2013	19/02/2014	19/03/2014
Subtidal	3/12/2013		

The intertidal¹, near shore mudflats were sampled at every sampling event. The sediment at these locations was exposed or submerged for no more than 50 cm. In addition, samples were taken from subtidal locations (in about 1 m water depth) at all ten sites in December 2013. The water level in the Murray Mouth and Coorong was very low during subtidal sampling, exposing a lot of tubeworm reefs in the Coorong, which made navigation difficult. Sampling for the subtidal locations took place at least 150 m towards the deeper channel from the intertidal locations. Two further intertidal mudflats on the Youngusband Peninsula were sampled in December 2013, to obtain two transects across the Coorong and Murray Mouth at Pelican Point and at Long Point. Unlike 2012/13, when subtidal and peninsula locations were sampled more frequently, they were only included in one of the sampling events in the 2013/14 monitoring.

At the intertidal sites, exposure varied on the three sampling occasions. With varying water levels in the Coorong and Murray Mouth, affected by tides, wind seiching and water releases, the widths of the mudflats ranged from being totally submerged to being exposed for >100 m. In late November, the

¹ Note the terminology is used although tides are only affecting the Murray Mouth region. Exposure of mudflats is also affected by variations in water from wind seiching, tidal prism and water release over barrages.

mudflats sampled in the Murray Mouth were exposed, yet during sampling in early December for the remaining sites, water levels were rising quickly and sampling took place in knee-deep water. During February 2013, water levels were again variable, even among sites within regions sampled on the same day. Wide mudflats were exposed at Ewe Island (200 m), Pelican Point (100 m) and Villa de Yumpa (250 m), but only narrow sections of mudflat were exposed at other sites (between 2 m at Long Point – 50 m at Parnka Point South). Water levels continued to fluctuate over summer and were generally high during March 2013, when only narrow areas of mudflat were exposed (between 2 m at Pelican Point – 100 m at Villa dei Yumpa) at the time of sampling.

Patches of *Ruppia spp.* (cf. *tuberosa* and *megacarpa*) were observed at most sites during December 2013, with some flowers observed on *R. cf. tuberosa*.

3.2 Environmental parameters

Water quality was measured in the field using a hand-held YSI Pro2030 multi-parameter electrode to record water temperature (° C), salinity (ppt), oxygen concentration (O₂ mg/l) and saturation (O₂ %). The pH of the water overlying the mudflats (intertidal) or subtidal sediments were measured using pH indicator-strips (Acilit 0-14).

Sediment characteristics were determined from samples taken in the field, frozen and later analysed in the laboratory. Three replicate samples were taken per site and location for each of the following variables to characterise sediment properties; grain size, organic matter and chlorophyll-a.

For sediment grain size, samples were taken with a 60 ml cut-off syringe (surface area 6.6 cm² to a depth of approximately 10 cm). Grain size was determined by laser diffraction using a particle size analyser (Malvern Mastersizer 2000). Sediment grain size samples were thawed and the fraction >1 mm sieved off manually to avoid blockage in the machine. The weights of this fraction and of the remaining sediment were determined for later normalisation of the data to correct for this procedure. Median and quartiles, as well as percentage of various particle sizes were obtained from the Mastersizer output. Sediment sorting (S_o) was calculated from the ratio of the quartiles (P₂₅ and P₇₅) $S_o = (P_{25}/P_{75})^{1/2}$, based on the metric scale.

For sediment organic matter, samples were taken with a 60 ml cut-off syringe (surface area 6.6 cm² to a depth of approximately 10 cm). Sediment samples were dried to constant weight using an Ohaus MB45 Moisture Balance. The sediment was homogeneously distributed onto aluminium trays and dried using the standard drying protocol (controlling the temperature profile at 80 °C). The profile burn was automatically completed after all moisture content was dried and remained stabilised for 30 seconds. Samples were then burnt in a muffle furnace at 450 °C for 5 hrs. Organic matter was thus determined by combustion and expressed as % dry weight (d.w.).

For sediment chlorophyll-a, samples were taken by inserting a 5 ml vial about 1 cm into the sediment. While in the field, 5 ml of methanol was added to extract the chlorophyll, and the vial was thoroughly shaken before being wrapped in aluminium foil (Seuront and Leterme 2006) and frozen for later analysis with a fluorometer (Turner 450). After the initial reading for total chlorophyll, drops of 0.1 M hydrochloric acid (HCl) were added to the samples to correct for phaeophorbides.

3.3 Macrofauna

Samples for macroinvertebrates in the sediments of the Murray Mouth and Coorong were taken with handheld PVC corers (83.32 cm² surface area). The corer was inserted into the sediment to approximately 15 cm depth, sealed with a stopper to avoid disturbance of the sample and dug out with a shovel or by hand. For subtidal locations, samples were taken using the same PVC corer, as the water level was very low (< 1 m) at the time of sampling, allowing operators to leave the boat. At each site and location, ten haphazardly placed replicate samples were taken for macroinvertebrates at each of the three sampling times.

Samples were sieved through a 500 µm mesh size while still in the field. In the laboratory, samples were sorted within several days while organisms were alive. Due to the sample and specimen volume for each sampling occasion, the sorted specimens were preserved in 70% ethanol for later identification. When not all samples could be processed within a few days, the entire sample was preserved in 70% ethanol. This occurred only in December 2013, when an unexpected high number of macroinvertebrates was encountered in the samples.

Identification was carried out under dissecting microscopes, and organisms were identified to the lowest possible taxonomic level and the number of individuals for each species per sample recorded. Amphipods were not differentiated into family or species. For insects, larval and pupae life stages were distinguished during sorting, but considered together for data analysis.

An assessment of recruitment of major macroinvertebrate species was carried out by size frequency analyses of the polychaete species *Capitella* sp., *Simplisetia aequisetis* and *Boccardiella limnicola* and the micro-mollusc *Arthritica helmsi*. When abundances of these species were high, no more than 30 individuals per species and sample were measured, if <30 individuals occurred in a sample, they were all measured. Too few specimens of *Australonereis ehlersi*, and *Nephtys australiensis*² were found for size-frequency analysis. Only complete specimens were used for size determinations, and length was measured on laminated graph paper under a dissecting microscope to the nearest 1 mm accuracy. Measurements were also taken of additional macroinvertebrates (e.g. bivalves) to obtain a range of their sizes.

3.4 Data analysis

To test for differences in salinity between the sampling sites, only intertidal locations sampled on all three occasions in the 2013/14 survey were considered, with month as factor (Dec-13, Feb-14 and Mar-14). Differences in salinity between the inter- and subtidal as well as peninsula locations were tested in December-13, the only months in the current survey when all locations were sampled. Data were square-root transformed before permutational analysis of variance (PERMANOVA) using a Euclidian distance-based similarity matrix.

²Note that *Nephtys australiensis* is now recognised as *Aglaophamus australiensis* but *N. australiensis* is retained throughout this report for ease of comparison to earlier reports and data sets.

To test for differences in the total abundance of macroinvertebrates in all survey years, data were fourth root transformed before calculating a Euclidean distance matrix (as single variable) used with a PERMANOVA design including year (2010/11, 2011/12, 2012/13, 2013/14) and region (Murray Mouth, North Lagoon, South Lagoon) as fixed factors.

To test for differences in total abundances between the depth locations (intertidal, subtidal and peninsula (at two sites)) for the survey in December 2013, data were prepared as above and a design including site and depth as fixed factors. Pairwise tests were run to assess site-specific differences in abundances with depth locations. To investigate recruitment of macroinvertebrate species and recolonisation of mudflats (key questions 1b-g), current (2013/2014) distribution ranges, abundances of key species (e.g. polychaetes) and taxonomic groups (e.g. amphipods, larval insects) as well as size frequency distributions were determined and compared to previous surveys.

Macroinvertebrate communities over the survey periods and study regions were characterised based on SIMPER analysis identifying species contributing most to communities within sites or time periods, or to the differentiation between sites, regions, survey years or periods. These species lists were compared with historical information (Geddes & Butler 1984; Geddes 1987) and records from earlier monitoring (Dittmann *et al.* 2010, Dittmann *et al.* 2011, Keuning *et al.* 2012, Dittmann *et al.* 2013a, b) and long-term data from TLM monitoring (Dittmann *et al.* 2014). For long-term community comparisons, including the TLM monitoring timeframe (Dittmann *et al.* 2014), the years were divided into three periods; pre-drought (2004-2006), drought (2007-2009), and flow (since 2010). To visualise differences in macroinvertebrate communities, Principle Coordinate Plots (PCO) were created, based on fourth root transformed data and Bray-Curtis similarity with a dummy value due to many zero values in the data. Canonical analysis of principle coordinates (CAP) discriminate analysis based on different flow scenarios was used to determine if distinct communities existed during different flow scenarios in each region. Samples collected during the 2013/2014 survey were then fitted to the CAP model generated for each location and assigned to a community type based on the two flow regimes to investigate the current state of macroinvertebrate communities within the Coorong.

Links between environmental conditions (using variables for water and sediment quality) and macroinvertebrate communities reestablishing in the Murray Mouth and Coorong were analysed through distance-based linear models (DistLM), visualised with distance-based redundancy analysis (dbRDA) plots. All analyses were carried out using the software PRIMER v6 with PERMANOVA add-on.

4. Results and Discussion

4.1 Environmental Conditions

Key Question 1a) Will environmental conditions within the Murray Mouth reflect true estuarine conditions and have they further improved in the North and South Lagoons since flows recommenced in 2010?

Characteristic of estuarine conditions include seasonal variations in salinities with riverine inflow or drought, and stratification of salinities resulting from freshwater inflow or saltwater intrusion (Potter et al. 2010). The salinity profiles in estuaries are one of the main determinants of estuarine fauna (Whitfield et al. 2012), and salinity had emerged in previous monitoring as the main variable affecting macroinvertebrate distributions in the Murray Mouth and Coorong (Dittmann et al. 2013a). During the 2013/14 response monitoring for macroinvertebrates, salinities measured during the sampling events gave indications for estuarine conditions improving.

In the December survey, salinity measurements were taken from water adjacent to intertidal mudflats (including two locations on the peninsula) and overlying subtidal sediments. Salinities were higher at most of the subtidal sites (Figure 3, Pseudo- $F = 101.41$, $P_{(perm)} = 0.0001$), yet measurements from two different water depths gave no indication of stratification. Salinities were also higher in the water adjacent to intertidal mudflats sampled on the peninsula locations of Pelican Point and Long Point compared to the mainland shore of the Coorong (Figure 3).




Figure 3. Salinity (average \pm SE) at each study site during sampling in December 2013 for the macroinvertebrate survey. Note that the intertidal and subtidal locations were not sampled on the same day. Sites are labelled as in Table 2, the suffix -P indicates locations on the peninsula.

The well-known salinity gradient, increasing from brackish-marine conditions in the Murray Mouth to hypersaline conditions in the Coorong, persisted in the current study period (Figures 3 and A1). At the three surveys over summer 2013/14, salinities changed significantly by site and date (Pseudo- $F = 473.75$, $P_{(perm)} = 0.0001$).

In the Murray Mouth, salinities were oligohaline (0.8 ppt (Monument Road) – 5.8 ppt (Hunters Creek)) at all four sites during December 2013, following higher spring flow (Figures 4 and 5). Yet higher salinities (up to 12.3 ppt) were recorded in water puddles on top of the mudflat at Ewe Island at low

tide. With reduced freshwater inflow over summer, salinities increased in the Murray Mouth and were meso-haline (24 ppt (Monument Road) - 31 ppt (Ewe Island)) in February, reaching marine salinities at Ewe Island in March (35 ppt) (Figure 4).

In the North Lagoon, salinities during December 2013 were mesohaline at the northern-most sites Mark Point (11 ppt) and Long Point (28 ppt), increasing to hypersaline at Noonameena (52 ppt) and Parnka Point (69 ppt, north of Hells Gate) (Figure 4). This wide range of salinity narrowed over the




Figure 4. Salinity (average \pm SE) at each study site for each sampling occasion during the 2013/2014 macroinvertebrate survey. Sites are labelled as in Table 2. The northern side of Parnka Point (PaPN) was only sampled in December 2013 in conjunction with The Living Murray condition monitoring. Note different scales of the y-axis. The dashed line shows the salinity of seawater (36 ppt) for comparison among panels.

following summer sampling events and seawater salinities (~36 ppt) were recorded in March 2014 at all sampled sites. Salinities had increased over summer at the northern section of the North Lagoon, but decreased at the southernmost sections (Figure 4).

In the South Lagoon, salinities remained hypersaline and less variation was recorded between the three sampling events. Parnka Point South (just south of Hells Gate), salinity was more stable, with a slight decrease in salinity from 75 ppt in December 2013 to 63 ppt in March 2014 (Figure 4). Salinity at Villa de Yumpa rose over the study period from 68 ppt in December 2013 to 93 ppt in March 2014.




Figure 5. Average salinity at those study sites in the Murray Mouth and Coorong surveyed for several summer surveys of the water release project since December 2010. Note that not all sites were sampled in every summer period. Sites in the Murray Mouth are Monument Road (MR), Hunters Creek (HC), Ewe Island (EI), Pelican Point (PP), and sites in the Coorong comprise Mark Point (MP), Long Point (LP), Noonameena (NM), Parnka Point (PaP) and Villa de Yumpa (VdY). Note different scales of the y-axis. Flow data are based on MDBA (see Figure 1).

Since flows resumed, salinities reduced substantially (by >30 ppt) in the Murray Mouth, North and South Lagoon (Appendix Figure A2) (Dittmann et al. 2014). The more frequent sampling over the summer months for the water release monitoring indicates fluctuations in salinity in response to the flow volume (Figure 5). The lower salinities in spring following freshwater inflow over winter, and the higher salinities over summer when flows were reduced, reflect estuarine conditions. This pattern has been apparent in the Murray Mouth over the last two summers and a similar seasonal variation in salinity can also be seen from the Coorong, especially at the sampling sites in the North Lagoon (Figure 5). A continuation of seasonal freshwater flows is needed to further improve the estuarine characteristics of the Murray Mouth and Coorong.

4.2 Macroinvertebrate Recruitment

Key Question 1b) Has recruitment continued for macroinvertebrate species that began recovery in 2011-12 and is recruitment occurring for other species in 2013-14?

Recolonisation of marine sediments disturbed on a large scale relies on larval influx from source populations elsewhere (Thrush et al. 1996, Whitlatch et al. 1998, Cummings & Thrush 2004). Larvae and juvenile stages have a higher dispersal potential than adult macroinvertebrates (Günther 1992), although emergence of larger benthic organisms has been documented (Oishi & Saigusa 1999). For the recolonisation of mudflats in the Murray Mouth and Coorong, presence of small sized individuals (assumed to have been recently recruited to the population) gives a good indication of the potential for recovery. Based on findings from the previous surveys we had predicted that recruitment of polychaetes and other benthic taxa would occur, and that additional species would recruit into the sediments. The analyses of size frequency distributions of polychaetes and records of size ranges in the summer survey 2013/14 indicated that several of the key macroinvertebrate species, such as of *Simplisetia aequisetis*, *Capitella* spp. and *Boccardiella limnicola* recruited throughout the summer months (Table 3).

The polychaete *Simplisetia aequisetis* ranged in size from 2 mm to 67 mm length (Appendix Table A1). Small-sized individuals were observed in the Murray Mouth and North Lagoon during late spring and early summer and, in the Murray Mouth, also in late summer (Figure 6). Size-frequency distributions for this species were similar in the Murray Mouth and North Lagoon regions during December 2013 and February 2014, while smaller and mid-sized worms were more frequent in the Murray Mouth than North Lagoon in March 2014. The presence of small individuals in each month indicates continuous recruitment over the survey period, yet there was little increase in the frequency of larger worms in the following months.

Size frequency distributions of *S. aequisetis* were similar between inter- and subtidal locations in the Murray Mouth region in December 2013, with a high proportion of small-sized individuals indicating recruitment in mudflats and shallow subtidal sediments (Figure 6). In the North Lagoon, the proportion of small-sized individuals was higher in the subtidal than intertidal mudflat samples (Figure 7). The size-frequency plots also showed that larger specimens of *S. aequisetis* were mostly found in the subtidal sediments (Figure 7).

Over the last three summer surveys, the size-frequency distributions of *S. aequisetis* were comparable and skewed to the left, with a higher frequency of small (~6-12 mm) than large worms (Appendix Figure A8).

Another polychaete related to *S. aequisetis*, *Australonereis ehlersi*, had been found in the previous surveys with a wider size range, indicating recruitment and large mature worms, occurring mainly in subtidal sediments (Appendix Figure A10 and Dittmann et al. 2013a). This species was rare in the recent survey as only one large individual of 138 mm length was found in the mudflat at Ewe Island.




Figure 6. Size-frequency distribution histograms for *Simplisetia aequisetis* for each sampling occasion in the Murray Mouth and North Lagoon over the summer 2013/14 survey. The number of individuals measured in each month is given by *n*.




Figure 7. Size-frequency distribution histograms for *Simplisetia aequisetis* for both subtidal and intertidal populations in a) the Murray Mouth and b) the North Lagoon sampled during December 2013. The number of individuals measured in each region is given by *n*.

During the 2012/13 survey, *A. ehlersi* was recorded almost exclusively at Noonameena and most abundant in the subtidal in the September 2012 survey (Dittmann *et al.* 2013a). The population size and structure of this species could have been underestimated in the present survey as subtidal sediments were only sampled once.

The polychaete *Capitella* sp. ranged in size from 2 mm to 18 mm length (Appendix Table A1). Large populations of this species were only found in the North Lagoon, at Long Point and Noonameena during December 2013, at Long Point in February 2014 and at Noonameena in March 2014 (Figure 8). The size frequency distributions gave no indication of any particular recruitment event that was captured in the samples, and worms of all sizes for this species were found in every month. The continued presence of small-sized individuals may indicate recruitment of this opportunistic species throughout the summer months.

Smaller worms occurred in samples from Noonameena compared to medium to large sized at Long Point. As Noonameena is located further south in the North Lagoon, this size-frequency distribution may hint to more recent colonisation at Noonameena. Specimens of *Capitella* sp. found in subtidal sediments of the North Lagoon in December 2013 were larger than in intertidal sediments (Figure 9). This distinct difference in the size frequency distribution could indicate recruitment or colonisation in the mudflats.

Size frequency distributions of *Capitella* sp. were similar over the last three summer surveys and unimodal, with a high frequency of worms 8-10 mm in lengths (Appendix Figure A9). Smaller specimens occurred each year, yet the frequency of larger specimens was more prominent in the previous two summers. In 2011/12 and 2013/14 the frequency of medium and large sized *Capitella* sp. appeared to be higher in subtidal sediments.




Figure 8. Size-frequency distribution histograms for *Capitella* sp. for Long Point (LP) and Nooneameena (NM) in the North Lagoon for each sampling occasion of the survey over summer 2013/14. The number of individuals measured in each month is given by *n*.




Figure 9. Size-frequency distribution histograms for *Capitella* sp. for both subtidal and intertidal populations in the North Lagoon sampled during December 2013. The number of individuals measured in each depth is given by *n*.

Few ($n=21$) specimens of *Nephtys australiensis* were found in early summer 2013/14 and their length ranged from 12 to 58 mm. Small-sized specimens occurred in subtidal sediments, and the two largest worms were found in the mudflat mainly at Ewe Island. As in previous surveys, no recruitment peaks were apparent from the size-frequency distributions, and worms of all sizes were present (Appendix Figure A11). The range of sizes measured in summer 2013/14 was narrower than the size range found in summer 2011/12 (Appendix Figure A11). With no further subtidal sampling beyond December 2013, it is unknown whether this species persisted in subtidal sediments.

The spionid polychaete, *Boccardiella limnicola*, ranged in size from 2 mm to 22 mm length (Appendix Table A2) and was only found at sites in the Murray Mouth. The size frequency distribution in December 2013 was bimodal, with a high frequency of small sized worms in both inter- and subtidal sediments indicating possible recent recruitment, and a second peak of larger worms around 15 mm lengths (Figure 10a). A similar recruitment event in late spring was also observed in the 2012/13 survey (Dittmann *et al.* 2013). By February 2014, worms of all sizes were found with similar frequency (Figure 10b). The sample size for size measurements of *B. limnicola* was small and only two individuals could be measured during March 2014, which were larger specimens (14 and 16 mm lengths).

Taking the size measurements of *B. limnicola* from over all survey months of the last two summer monitoring periods, a spread of smaller and larger worms was found in each year, slightly skewed to smaller sized specimens, especially in the subtidal when taken over the entire summer 2013/14 (Appendix Figure A12).




Figure 10. Size-frequency distribution histograms for *Boccardiella limnicola* for a) both subtidal and intertidal populations in the Murray Mouth sampled during December 2013, and intertidal populations in b) February 2014. Only two specimens were found in March 2014 of 14 and 16 mm lengths. The number of individuals measured in each month is given by *n*.

Table 3. Occurrence of small individuals of key species observed in benthic samples on each sampling occasion. Subtidal sampling was only conducted during December 2013. Ticks indicate small individuals were present at the specified location on the sampling occasion.

Species	Location	Small individuals observed			
		Dec-13		Feb-14	
		Intertidal	Subtidal	Intertidal	Intertidal
<i>Simplisetia aequisetis</i>	Murray Mouth	✓	✓	✓	✓
	North Lagoon	✓	✓	✓	✓
<i>Capitella spp.</i>	Murray Mouth				
	North Lagoon	✓	✓	✓	✓
<i>Boccardiella limnicola</i>	Murray Mouth	✓	✓	✓	
	North Lagoon				
<i>Arthritica helmsi</i>	Murray Mouth	✓	✓	✓	✓
	North Lagoon	✓	✓	✓	

The micro-mollusc *Arthritica helmsi* is a small sized bivalve which had been prominent in the Murray Mouth and Coorong before the drought and started to return last year (Dittmann et al. 2013 a, b). The population size of this species was higher again in the recent survey and they were found at all three sampling events. The size-frequency distributions were uni-modal with most bivalves 2 mm in size

(Figure 11). The sizes recorded for this species ranged from 0.5 to 4 mm (Appendix Table A2). The presence of juveniles is difficult to determine due to the small size of this species. Yet, individuals <1 mm in length were observed on each sampling occasion, which may represent recent recruitment. Size frequency distributions of specimens collected from inter- and subtidal locations were similar in December 2013 (Figure 12).




Figure 11. Size-frequency distribution histograms for *Arthritica helmsi* for each sampling occasion in the Murray Mouth and North Lagoon regions.




Figure 12. Size-frequency distribution histograms for *Arthritica helmsi* for both subtidal and intertidal populations in the Murray Mouth region sampled during December 2013.

Amphipods were abundant in the December 2013 survey, and females carrying eggs were frequently observed. Too few complete specimens of other macroinvertebrate species were collected to obtain detailed size-frequency distributions and assessment whether recruitment occurred. The size ranges of macroinvertebrates encountered are provided in Appendix Table A2.

Insect larvae, in particular chironomids (non-biting midges), have been a prominent component of the macroinvertebrates in the Murray Mouth and Coorong, particularly in the South Lagoon (Geddes and Butler 1984, Dittmann et al. 2013a). This pattern continues with high abundances of chironomids in inter- and subtidal sediments north of Paraka Point and at Villa de Yumpa (Figure 6a). Chironomidae larvae were most abundant during December 2013, and found in the Murray Mouth and Coorong. Larval stages of Ephydriidae (brine fly) and Dolichopodidae (long-legged flies) were observed during all sampling occasions and at most of the sampling sites (Figure 13b and c). Other insect larvae were occasionally recorded in samples from a few sites, such as Muscidae (house flies) and Coleoptera (beetle larvae) in March 2014 only, and Ecnomidae (caddisfly larvae) during February and March 2014 (Figure 13d-f). The presence of insect larvae in sediments and waters of the Murray Mouth and Coorong is subject to their life history and thus rather ephemeral.

Key Question 1c) Can the recruitment of species in 2013/14 be linked with the timing of differing flow scenarios to identify drivers of macroinvertebrate recruitment?

The comparison of flow data and salinity (Figure 5) indicated a return of seasonal estuarine salinity variation, which can affect recruitment of estuarine organisms (Gillet and Torresani 2003). Juveniles of several key species were found at the sampling events in December 2013, February 2014 and, to a lesser extent, in March 2014 (Table 3). The reduced flow volume and increasing salinities in early summer (Figure 5) may provide a trigger for recruitment of some species, such as *Boccardiella limnicola* and possibly the crab *Paragrapsus gaimardii* and mysid shrimps which were encountered in samples from December only (Appendix Table A1). Settlement of larvae can be triggered by cues




Figure 13. Average abundance of larval insects observed in samples at each study site in the Murray Mouth and Coorong during the 2013/14 survey period. In December 2013, inter- (suffix –int.) and subtidal (suffix –sub.) locations were sampled. All other locations were intertidal. The peninsula locations at Pelican Point (PP-p) and at Long Point (LP-p) were only sampled in December 2013. Error bars indicate standard deviation.

other than water quality, as sediment characteristics and biotic interactions can further affect settlement and thus recruitment patterns of macroinvertebrates (Olafsson et al. 1994, Sebasvari et al. 2006).

Juvenile bivalves and *Nephtys australiensis* were mainly encountered in the December sampling event (Appendix Table A1). As subtidal sampling was carried out only once in December, recruitment in subtidal sediments at other times over the summer survey 2013/14 may have been missed.

Some of the key macroinvertebrate species occurred with a wide size range at each sampling event over summer 2013/14, which could indicate adaptation to estuarine conditions and continuous recruitment independent of environmental triggers. The population biology of key species such as *Simplisetia equisetis* would thus be of interest to be explored further.

4.3 Macroinvertebrate Recolonisation

Key Question 1d) Will there be any further recolonisation of mudflats by macroinvertebrates absent from areas in previous monitoring 2010-12?

At the three sampling events in summer 2013/14, 32 macroinvertebrate taxa were differentiated in the samples, composed of molluscs (12 taxa), annelids (8 taxa), insect larvae (7 taxa) and crustaceans (5 taxa) (Appendix Table A1 and Figure A7). The total number of species found in intertidal mudflats had thus increased over previous monitoring (2010/11, 2011/12 and 2012/13: 14, 21 and 20 taxa respectively (Dittmann et al. 2011, Keuning et al. 2012, Dittmann et al. 2013a)). Looking at species numbers by region, the further increase was less pronounced (Figure 14). As sampling effort varied over the monitoring years (number of sampling events, sites and locations sampled), species numbers for the Coorong and peninsula locations have to be considered with caution. The number of species found in mudflat samples also varied within sites over the three sampling events in 2013/14, without any consistent pattern of increase or decrease (Appendix Figure A6).

Two species were recorded in the 2013/14 survey for the first time again since the drought and flow resumed, the scavenging polychaete *Phyllodoce novaehollandiae* at Ewe Island and an isopod at Parnka Point. Juvenile bivalves of possibly three different species were found in the December sampling event. Larvae of Ecnomidae and Coleoptera had not been found in the previous water release monitoring.

The increase in the total species number in 2013/14 was largely due to molluscs. Six morphospecies of small hydrobiid snails were identified, which occurred at sites in the Murray Mouth. Hydrobiids were rarely encountered in previous surveys and species not consistently differentiated. For a comparison of species numbers over the survey years, they were thus combined to family level (Hydrobiidae) and the real species numbers in 2013/14 will be higher than the comparison shown in Figure 14.

For mudflat sediments, abundances of macroinvertebrates in the current monitoring year were comparable or lower than in the previous surveys since flow resumed, while similar or higher abundances were recorded for the subtidal study sites (Appendix Figure A13). Abundances of macroinvertebrates in mudflat sediments were highly variable within each of the survey years, in particular in the Murray Mouth, where a decrease was recorded over summer (Figures 14 and 15, Appendix Figure A14). Abundances were significantly different between the survey years (Appendix Table A3). Yet for intertidal mudflats in the Murray Mouth and North Lagoon, abundances in 2013/14 were not significantly different to those found in 2011/12, possibly affected by the decline over summer 2013/14. For the South Lagoon (note, only the northern end to Villa de Yumpa sampled since 2011/12), macroinvertebrate abundances in mudflats were significantly higher in the 2013/14 survey than in any of the previous two survey years. In mudflats at the two peninsula locations first sampled in 2012/13, abundances were similar between years for Pelican Point in the Murray Mouth, but had significantly ($P_{(perm)}=0.0001$) increased at the peninsula mudflat at Long Point by 2013/14 (Figure 14). For macroinvertebrate abundances in subtidal sediments, pairwise tests showed significant increases between any two years for the three regions (Appendix Table A3).




Figure 14. Average abundance (individuals per m²; with standard deviation) of macroinvertebrates and total number of species recorded (Total S) in samples from the monitoring periods since flow resumed in late 2010. a) the Murray Mouth, b) the North and c) South Lagoon regions. Sampling periods for the water release monitoring were 2010/11: December 2010 – April 2011; 2011/12: October 2011 – May 2012; 2012/13: September 2012 – March 2013; 2013/14: December 2013 – March 2014 = this current report). 2010/11 and 2011/12 include the Tauwicheere intertidal site on the peninsula. Asterisks indicate that no sampling occurred for CLLMM monitoring in the survey period and region. Note that different sites and different numbers of sites were included for the Murray Mouth and North Lagoon regions for each monitoring time span, and that subtidal and peninsula locations were only sampled during December 2013 for the 2013/14 monitoring.

Assessing further recolonisation of mudflats in 2013/14 over previous surveys was affected by a pronounced decrease in abundances over the summer months (Figure 15). At all of the Murray Mouth sites and most of the sites in the Coorong, abundances had dropped several orders of magnitude between December and February, and remained low into March. In the Coorong, the decrease was particularly pronounced at Noonameena, where abundances had dropped >65x since December, while macroinvertebrate abundances at Long Point and Parnka Point South changed less over the three sampling events (Figure 15).

A further indication of incomplete recolonisation in mudflats became apparent in December 2013, as abundances were only significantly higher in intertidal than subtidal locations at two sites (Hunters Creek and Ewe Island) whereas at all other sites, abundances in subtidal locations significantly

exceeded those found in intertidal mudflat sediments (Figure 15, Appendix Table A4). At the two sites where a transect was sampled from land-based mudflat over the subtidal to peninsula mudflat, abundances did not differ between the two mudflats at Pelican Point, whereas at Long Point, abundances were significantly higher at the peninsula mudflat (Figure 15, Appendix Table A4).

The differences in abundance between inter- and subtidal sediments in December 2013 were driven by the prominent macroinvertebrate species occurring in the Murray Mouth and Coorong (Figure 16). Amphipoda were more abundant in mudflats than subtidal sediment at Hunters Creek and Ewe Island and accounted for most of the overall pattern in abundance between the inter-, subtidal and peninsula locations in December 2013 (Figures 15 and 16). Chironomid larvae and the polychaete *Simplisetia aequisetis* contributed to the higher abundance in mudflats at Hunters Creek and Ewe Island as well (Figure 16).




Figure 15: Average abundances (individuals per m²; n=10, with standard deviation) of total macroinvertebrates recorded at intertidal, subtidal and peninsula locations during December 2013, and at intertidal locations sampled in February and March 2014. See Figure 2 for site locations.















Figure 16 (previous page). Average abundances (individuals per m²; *n*=10, with standard deviation) of several abundant macroinvertebrate taxa recorded at intertidal, subtidal and peninsula locations during December 2013. Note the differences in the scale of the y-axis. See Figure 2 for site locations. The high error bars are a reflection of the patchiness of benthic macroinvertebrates.

For the numerically dominant amphipods, abundances in intertidal mudflats decreased from Pelican Point into the Coorong, while they were abundant in subtidal sediments. Abundances of amphipods collapsed over summer (Figure 17). The polychaete *Simplisetia aequisetis* had a similar distribution range as the amphipods, and was abundant in mudflats at most of the Murray Mouth and North Lagoon at all three sampling events (Figures 16 and 17). Both amphipods and *S. aequisetis* were, however, constricted to subtidal sediments at Mark Point in December 2013 and continued to be absent from mudflats at this sites in February and March 2014, yet occurred in mudflats further south in the Coorong at Long Point (Figures 16 and 17). Another polychaete, *Nephtys australiensis*, which was common in the past, was recorded from Hunters Creek and Ewe Island mudflats only, and subtidal sediments at Ewe Island.

The high abundance in mudflats on the peninsula location at Pelican Point was composed of *S. aequisetis* and the small bivalve *Arthritica helmsi*, whereas at Long Point, amphipods and the polychaetes *Capitella* sp. and *S. aequisetis* contributed most (Figure 16). At Long Point and Noonameena, *Capitella* sp. occurred in similar abundances in inter- and subtidal sediments and, at Long Point, also on the peninsula mudflat (Figure 16). This pattern was similar to the occurrence of *Capitella* seen in previous surveys (Dittmann et al. 2013a), yet this year the species was also found in mudflats and subtidal sediments at Parnka Point.

Ostracoda and larvae of Ephydriidae occurred almost exclusively at mudflats in the Coorong, with ostracods very abundant in sediments on the southern side of Parnka Point, accounting for the high overall abundance at this site (Figures 15 and 16). Chironomid larvae were also abundant at Parnka Point and in the South Lagoon, yet found mostly in subtidal sediments (Figure 16).

The potential for further recolonisation of intertidal mudflats arises from the presence and abundance of several macroinvertebrate species in sediments of adjacent subtidal sampling locations (Figure 16). For the bivalves *Soletellina alba* and *Spisula trigonella*, specimens were found mostly or only in subtidal sediments. At Hunters Creek, *S. trigonella* occurred in subtidal sediments in December 2013 and was found in mudflat sediments in February 2014 (Figure 17). Mysid shrimps and also the snail *Hydrobia* sp. 2 were more numerous in subtidal sediments at some sites in the Murray Mouth. *Arthritica helmsi*, which was prominent in the system before the drought and started to reappear in 2012/13 (Dittmann et al. 2007, 2013b) also re-established in mudflats throughout most of the Murray Mouth and into the North Lagoon over this survey (Figure 17).

As subtidal sediments were only sampled in December 2013, it is unknown whether the decrease in abundances in mudflats over the summer months occurred at these locations as well. In previous monitoring, indications were found that mudflats were recolonised from subtidal locations (Dittmann et al. 2013b). The size of the potential source populations of the named species for recolonising the mudflats could thus not be assessed in the 2013/14 monitoring.




Figure 17 (previous page). Average abundances (individuals per m⁻²; n=10, with standard deviation) of several abundant macroinvertebrate taxa recorded at intertidal locations during December 2013, February and March 2014. See Figure 2 for site locations. The high error bars are a reflection of the patchiness of benthic macroinvertebrates.

Key Question 1e) Will recolonisation of macroinvertebrate species occur in the South Lagoon if salinities remain lowered within this region?

Prior to environmental flows, macroinvertebrate communities at the South Lagoon sites Parnka Point South and Villa de Yumpa were poor in species and abundances. Salinity is still hypersaline and has remained around ~70 ppt for the third survey period in a row, with some flow-related fluctuations over spring and summer (Figure 4). The macroinvertebrates present are mainly composed of halophylic species, specifically larvae of salt-tolerant Diptera (Chironomidae and Ephydriidae), oligochaetes, ostracods and amphipodes. Ostracods were particularly abundant in mudflats on the southern side of Parnka Point (Figures 16 and 17), and have not occurred in these numbers before. They accounted for the increase in abundances for the South Lagoon (Figure 14). In December 2013, single specimens of estuarine macroinvertebrates were found in intertidal and subtidal sediment samples from Parnka Point South and Villa de Yumpa. This includes a few individuals of the polychaete *Capitella* sp. and the bivalve *Arthritica helmsi*. *Capitella* sp. was already recorded at this site during the 2012/13 survey, also with few individuals only (Dittmann *et al.* 2013a). *Arthritica helmsi* has only been recorded from the South Lagoon with single specimens in 2006 (Dittmann & Nelson 2007) and 2013 (Dittmann *et al.* 2013a). The polychaete *Australonereis ehlersi* was found in subtidal sediments just north of Parnka Point. The presence of only a few individuals of these estuarine macroinvertebrates around Hells Gate and at Villa de Yumpa does not yet indicate that recolonisation of the South Lagoon has occurred, but that environmental conditions allow specimens to be present in this region from where they may lead to future recolonisation.

Key Question 1f) Have species maintained or further increased their distribution range in comparison to 2011/2012?

It was predicted that with progressively improving environmental conditions in the estuary and Coorong lagoon, the distribution of macroinvertebrate taxa will extend across multiple sites in the system. In December 2013, species numbers were higher throughout the Murray Mouth and at several sites in the Coorong than in previous surveys and abundances were high across the entire sampling area, especially in subtidal locations (Figure 18). At the mudflats sampled in February and March 2014, species numbers remained higher than during the early post-flow periods in the Murray Mouth, while abundances of macroinvertebrates decreased (Figure 18).

During the 2013/14 survey, the six most prominent macroinvertebrate taxa (Amphipoda, chironomid larvae, *Capitella* sp., *Simplisetia aequisetis*, *Australonereis ehlersi* and *Nephtys australiensis*) occurred at sites where they were found before in the post-flow periods, yet there were shifts in their abundances across the sites (Figure 19). They maintained their distribution range but, apart from *Capitella* sp., did not increase it.

Amphipoda were less abundant in the Coorong than in the 2012/13 survey, and their abundances in the Murray Mouth decreased over summer 2013/14, as it did in summer 2011/12 (Figure 19). In the previous survey, amphipod abundances were also lower in mudflats of the Murray Mouth in March 2013, but still high in subtidal sediments. As subtidal sediments were not sampled in February and March 2014, it is unknown whether abundances in the subtidal had decreased as well or remained high.

For chironomid larvae, a more disjunct distribution emerged, with higher abundances at the Murray Mouth sites and in the Coorong from Nooan southward, where chironomids were particularly abundant in subtidal sediments at Parnka Point and Villa de Yumpa (Figures 16 and 17). Chironomid larvae were not found in mudflats at Mark Point and also absent from Long Point in February and March 2014, when their abundance had dropped in general (Figures 17 and 19).




Figure 18. Patterns of change in the distribution of macroinvertebrates since flow commenced, illustrated by average abundances (individuals per m²) and number of species found in sediments of the Murray Mouth and Coorong sites during the water release study. Abundances and species numbers are indicated by colour codes (see legends). Sampling locations are differentiated as intertidal (i) and subtidal (s). Peninsula locations are not included in this comparison. Discontinued sampling sites in the Murray Mouth from the 2010/2011 and 2011/2012 monitoring periods are also omitted. For Parnka Point, the northern and southern side were considered together when both were sampled. Fields that are crossed out indicate that no sampling took place and the site or depth location. For Dec-10, data are used from the 2010/2011 TLM monitoring and an additional sampling event at Long Point.

The distribution range of *S. aequisetis* was similar to previous surveys, and this polychaete species has well established in mudflats in the Murray Mouth and at Long Point (Figure 19). At Mark Point, where abundances of *S. aequisetis* were high in 2012/13, very few or no specimens were found in mudflats in the 2013/14 survey. Yet, the December 2013 sampling showed that *S. aequisetis* was present with high abundances in the subtidal location at this site and even at Noonameena (Figures 16 and 19). Previously, specimens of *S. aequisetis* were found in subtidal sediments at Villa de Yumpa in February 2013, yet as no further subtidal sampling occurred in the present survey, any southward expansion could not be assessed.

The small polychaete *Capitella* sp., which is an indicator for polluted or eutrophic sediments (Tsutsumi 1990, Ramskov and Forbes 2008), was highly abundant in inter- and subtidal sediments in the mid-region of the North Lagoon (Long Point and Noonameena) during December 2013, where it occurred with consistently high abundances since flows resumed (Figure 16 and 19). The distribution range of this species was extending both towards the South Lagoon, with individuals found at Parnka Point and also subtidally at Villa de Yumpa in December 2013, and northward, where specimens were recorded in mudflats of Mark Point and the Murray Mouth sites in February and March 2014 (Figures 16 and 19). This coincided with an increase in salinities in the Murray Mouth (Figure 4) and an increase in sediment organic matter (Appendix Figure A4). In early 2014, the abundance of *Capitella* sp. decreased at Long Point and Noonameena, and this species was not recorded in mudflats south of Noonameena, similar to the March surveys in previous years (Figure 19).

Two other polychaete species, *Australonereis ehlersi* and *Nephtys australiensis*, occurred at very few sites in previous surveys and the current sampling (Figure 19). They were found at the same sites where they were recorded before and their numbers remained low, with no records from samples in March 2014. *Australonereis ehlersi* was only recorded in low abundances from mudflats at Noonameena and Ewe Island during the 2013/14 monitoring, and from subtidal sediments at Parnka Point (Figure 19). While specimens of this species had been recorded from the subtidal at Long Point in previous surveys, it was not found at this site in 2013/14. *Nephtys australiensis* maintained a small population at Ewe Island, and was also observed in the intertidal at Hunters Creek, where it occurred in subtidal sediments in February 2013 (Figure 19).

There were no indications for extensions of distribution ranges for further macroinvertebrate species in the 2013/14 monitoring period (Figure 20). Based on the number of sites of the continuously surveyed mudflats and subtidal sediments, distribution ranges had increased within one to two years after the high flow period of 2010/11, but have since been maintained or declined (e.g. for oligochaeta) (Figure 20). Some taxa have a wider distribution range in subtidal sediments (e.g. *Soletellina alba*), while others occurred at more sites in the intertidal mudflats (e.g. Ephydriidae). The assessment of distribution ranges is thus also affected by changes in sampling design between monitoring periods. Given the patchiness of several macroinvertebrate species on a small scale (within sites) (Figures 16 and 17), sample size can further affect the assessment of distribution and abundance. Figure 20 illustrates the presence of species at sites only, but not the abundance in their distribution range.

Amphipoda											Chironomidae											
		MR	HC	EI	PP	MP	LP	NM	PaP	VdY			MR	HC	EI	PP	MP	LP	NM	PaP	VdY	
Dec-10	i																					
	s																					
Feb-11	i																					
	s																					
Mar-11	i																					
	s																					
Apr-11	i																					
	s																					
Oct-11	i																					
	s																					
Dec-11	i																					
	s																					
Feb-12	i																					
	s																					
Mar-12	i																					
	s																					
May-12	i																					
	s																					
Sep-12	i																					
	s																					
Dec-12	i																					
	s																					
Feb-13	i																					
	s																					
Mar-13	i																					
	s																					
Dec-13	i																					
	s																					
Feb-14	i																					
	s																					
Mar-14	i																					
	s																					
<i>Simplisetia aequisetis</i>											<i>Capitella sp.</i>											
		MR	HC	EI	PP	MP	LP	NM	PaP	VdY			MR	HC	EI	PP	MP	LP	NM	PaP	VdY	
Dec-10	i																					
	s																					
Feb-11	i																					
	s																					
Mar-11	i																					
	s																					
Apr-11	i																					
	s																					
Oct-11	i																					
	s																					
Dec-11	i																					
	s																					
Feb-12	i																					
	s																					
Mar-12	i																					
	s																					
May-12	i																					
	s																					
Sep-12	i																					
	s																					
Dec-12	i																					
	s																					
Feb-13	i																					
	s																					
Mar-13	i																					
	s																					
Dec-13	i																					
	s																					
Feb-14	i																					
	s																					
Mar-14	i																					
	s																					

cont. next page




Figure 19. Changes in the distribution ranges and average abundances (individuals per m²) of key macroinvertebrate taxa, indicated by colour codes based on a log scale (see legend). Sampling locations are differentiated as intertidal (i) and subtidal (s). Peninsula locations are not included in this comparison as first sampled in 2012-2013. Discontinued sampling sites in the Murray Mouth from the 2010-2011 and 2011-2012 monitoring periods are also left out. Sites in the Coorong that are crossed out were not sampled in the earlier monitoring periods yet for Dec-10, data are used from the 2010-2011 TLM monitoring and additional sampling at Long Point. No subtidal sampling occurred in February and March 2014.

The bivalve *Arthritica helmsi*, which was not found for several years during the drought and after flow resumed, was present again at several sites in the Murray Mouth and Coorong since the 2012/13 monitoring period, and while this species did not extend its distribution range in 2013/14, abundances were higher at all sites where it occurred. In 2012/13, abundances were highest at Mark Point (Dittmann et al. 2013a), which shifted to Hunters Creek and the peninsula mudflat at Pelican Point in 2013/14 (Figures 16 and 17). A few individuals of *A. helmsi* were even found at Villa de Yumpa in the South Lagoon in December 2013 (Figure 19).

Another bivalve, *Soletellina alba*, had increased its distribution range to include most sites between Monument Road and Noonameena in March 2013 (Dittmann et al. 2013a), and was found again at Monument Road, Ewe Island and Long Point in 2013/14, but in lower numbers (Figure 17). The bivalve *Spisula trigonella* occurred in subtidal sediments at a few sites in the 2012/13 monitoring and was most abundant at Noonameena (Dittmann et al. 2013a), where it was found again in December 2013. In the 2013/14 monitoring, this species was also recorded from inter- and subtidal mudflats at Hunters Creek and the mudflat at Ewe Island (Figures 16 and 17). As subtidal locations were sampled less frequently in 2013/14, it is impossible to determine if the distribution range of these bivalves has changed during 2013/14.

Key Question 1g) Has the macroinvertebrate community been restored to pre-drought conditions following the 2011/2012 barrage flows into the Coorong?

Macroinvertebrate communities have changed away from those present during the drought and indications for a recovery were apparent. An analysis using pre-drought, drought and flow periods revealed an increasing differentiation of the communities currently present in the Murray Mouth and Coorong to the ones in the drought period (Figure 22). The communities in the flow period were also distinct to the pre-drought period.

In the Murray Mouth, macroinvertebrate communities grouped with those from recent monitoring events and became distinct to the drought years as well as 2010/11, when the high flow volume passed through the estuary (Figures 1 and 22). The community change over the last decade almost follows a circular path, and a trajectory indicated increasing similarity with pre-drought communities (Figure 21). There was little differentiation by sites in the Murray Mouth over the years (Figure 22), as all sampling sites were in the vicinity of barrages and similarly influenced by water releases creating more estuarine conditions.

Over the summer of 2013/14, communities in the Murray Mouth were characterised by the presence of Amphipoda, the polychaetes *Simplisetia aequisetis* and *Boccardiella limnicola*, chironomid larvae and the small bivalve *Arthritica helmsi* (Table 4). Further species present in low numbers were the polychaetes *Australonereis ehlersi* and *Nephtys australiensis*, as well as hydrobiid snails and the bivalves *Spisula trigonella* and *Soletellina alba* (Appendix Table A1). These species were also characteristic for communities under freshwater or estuarine conditions in the Coorong as described three decades ago by Geddes and Butler (1984) and Geddes (1987) (Table 5).




Figure 20. Occurrence by the number of sites where macroinvertebrate taxa were recorded in the Murray Mouth and Coorong over the monitoring period for the water release study. Only sites continued throughout the monitoring periods were considered, being five in 2010/11 and nine in the following periods. Note that the number of sampling events in each monitoring period varied between three and five surveys per summer period, and that subtidal locations were less frequently sampled than intertidal sites.

For the Coorong lagoons, differences in macroinvertebrate communities over drought and flow periods were less clear, with some overlap between communities during the drought and flow (Figure 21). The northern and southern sections of the North Lagoon had distinct macroinvertebrate communities, which each changed slightly over time and site specific responses. The macroinvertebrate community at Long Point, for example, changed little over the last ten years, while at Noonameena, large differences occurred between drought and flow periods (Figure 22). For the North Lagoon, a trajectory indicated that macroinvertebrate community was very different from all other years in 2010 when flow commenced, with only slight indication of a return in the last three years (Figure 21).

Over the summer of 2013/14, communities at sites in the northern part of the North Lagoon (Mark Point and Long Point) were characterised by the presence of *Capitella* sp., *Simplisetia aequisetis*, Amphipoda, Ephydriidae larvae and *Arthritica helmsi* (Table 4). This community was similar to those described historically as characteristic of estuarine to hypermarine communities in the Coorong (Table 5). Communities in the southern part of the North Lagoon (Noonameena and Parnka Point North) were characterised by the presence of larvae of salt tolerant Chironomidae and Ephydriidae, and the polychaete *Capitella* sp. (Table 4). In addition, Ostracoda were also recorded at Parnka Point North (Appendix Table A2). This community resembled characteristics of historical hypermarine to hypersaline conditions (Table 5). Communities in the South Lagoon were characterised by the presence of Ostracoda and chironomid larvae (Table 4). Larvae of Ephydriidae were also recorded in samples from both South Lagoon sites (Appendix Table A1). This community was characteristic of historically hypersaline communities (Table 5).

Table 4. SIMPER results showing average similarities (Ave. Sim.) and characteristic taxa that contributed highly (greater than 10%) to average similarity among samples within each region (divided into northern and southern reaches for the North Lagoon based on observed differences in salinities, see Figure 4), for each sampling occasion during the 2013/14 monitoring period. Taxa are listed in decreasing order of contribution to average similarity (percent contribution for each species is given in brackets). Site acronyms are listed in Table 2.

Region	Site	Dec-13		Feb-14		Mar-14	
		Ave. Sim.	Characteristic taxa	Ave. Sim.	Characteristic taxa	Ave. Sim.	Characteristic taxa
Murray Mouth	All	73.84	Amphipoda (44.7) <i>Simplisetia aequisetis</i> (24.0) Chironomidae larvae (16.7)	45.45	<i>Simplisetia aequisetis</i> (54.5) <i>Arthritica helmsi</i> (13.1) <i>Boccardiella limnicola</i> (10.2)	42.24	<i>Simplisetia aequisetis</i> (57.2) Amphipoda (13.6)
North Lagoon	Northern MP, LP	40	<i>Capitella</i> spp. (24.1) <i>Simplisetia aequisetis</i> (21.7) Amphipoda (20.3) Ephydriidae larvae (17.6) <i>Arthritica helmsi</i> (14.2)	25.74	<i>Capitella</i> spp. (43.6) <i>Simplisetia aequisetis</i> (26.7) <i>Arthritica helmsi</i> (16.3)	10.06	Ephydriidae larvae (70.3) <i>Simplisetia aequisetis</i> (15.9)
	Southern NM, PaPN	55.14	<i>Capitella</i> spp. (53.4) Chironomidae larvae (27.4)	13.15	Ephydriidae larvae (90.8)	25.59	Ephydriidae larvae (71.3) <i>Capitella</i> spp. (19.4)
South Lagoon	All	43.48	Ostracoda (74.6) Chironomidae larvae (24.6)	23.67	Ostracoda (95.6)	13.48	Ostracoda (73.4) Chironomidae larvae (25.2)

Table 5. Presence/absence of taxa from communities within the Coorong from historical sources (Geddes & Bulter 1984; Geddes 1987) and the current monitoring period 2013/14 (this report).

Taxa	Historical communities (Geddes and Butler 1984; Geddes 1987)				Current communities (this report)			
	Freshwater (0 - 2 ppt)	Estuarine (5 - 30 ppt)	Hypermarine (35 - 50 ppt)	Hypersaline (50+ ppt)	Murray Mouth	Northern North Lagoon	Southern North Lagoon	South Lagoon
<i>Simplisetia aequisetis</i>	absent	abundant			abundant	abundant		
<i>Australonereis ehlersi</i>	rare	present			rare	present		
<i>Nephtys australiensis</i>	absent	present			present			
<i>Ficopomatus enigmaticus</i>		abundant			rare			
<i>Capitella</i> sp.	absent		present		present	present	abundant	
<i>Arthritica helmsi</i>		present			abundant	abundant		rare
<i>Spisula</i> sp.	rare	present			present			
Hydrobiidae		abundant	present		present			
<i>Salinator fragilis</i>	rare		present		present			
Ostracoda				present			present	abundant
Isopoda				present			rare	
Amphipoda		abundant	abundant		abundant	abundant	present	rare
Chironomidae Larvae			present	present	abundant	rare	abundant	present
Ephydriidae Larvae			present	present	rare	abundant	abundant	present




Figure 21: Trajectories of community change in the Murray Mouth and North Lagoon, over monitoring years since 2004 (TLM data, Dittmann et al. 2014), based on average data per regions and year from surveys carried out in late spring/early summer. The trajectory line is connecting consecutive years.




Figure 22. PCO plots of macroinvertebrate communities in the Murray Mouth and Coorong, based on sites sampled since December 2004 until December 2012, including Monument Road, Ewe Island and Pelican Point for the Murray Mouth and Long Point, Noonameena, Parnka Point and Villa de Yumpa for the Coorong. Only sampling dates between October and December were considered to avoid any confounding seasonal effects. Data from 2004 to 2010 (Coorong data for 2010 only) are taken from annual TLM monitoring, and for 2010-2013/14 from water release monitoring. Please note that for each region, left and right panels show the same plot with symbols in the second panel showing differences among sites.

4.4 Relationships between Macroinvertebrate Communities and Environmental Conditions

Key Question 2) Which environmental conditions influence the distribution, abundance and community structure of macroinvertebrates?

Macroinvertebrate communities in the Murray Mouth and Coorong have been shown to be affected by water quality variables, in particular salinity with a pronounced gradient from the estuary to the lagoon (Appendix Figure A1), and sediment properties such as grain size composition, organic matter and microphytobenthic biomass (Dittmann et al. 2013a, b). The tight relationship between salinities and abundance as well as diversity of macroinvertebrates was evident from the data in the 2013/14 monitoring, as both abundance and species numbers decreased with increasing salinity (Figure 23). This pattern followed established relationships between salinity and estuarine macroinvertebrates (Attrill and Rundle 2002, Whitfield et al. 2012). Salinities in the Murray Mouth and Coorong have changed over time with drought and flow periods (Kangas & Geddes 1984; Geddes 1987; Dittmann et al. 2013b), and corresponding changes in macroinvertebrate communities were recorded through the long-term TLM condition monitoring (Dittmann et al. 2013b) and water release intervention monitoring (this report)(Figure 22).




Figure 23. Scatter plot of a) total benthos abundance and b) total number of species (macroinvertebrate taxa) versus salinity (x-axis). Each point represents a sampling location and occasion (pooled across replicates) during the 2013/2014 monitoring

To explore which environmental conditions could have affected the macroinvertebrate communities in the recent monitoring period, multivariate distance-based linear models (DistLM) with distance-based redundancy analysis (dbRDA) were carried out (Figure 24). These analyses revealed that the change in macroinvertebrate communities of the Murray Mouth and Coorong was again mainly driven by the higher salinity towards the South Lagoon (Pseudo- $F = 7.900$; $P_{(perm)} = 0.0001$), but the fine sand component of sediments affected macroinvertebrates as well (Pseudo- $F = 2.215$; $P_{(perm)} = 0.0388$). Yet, salinity and fine sand together explained only about 30% of the total variation in macroinvertebrate community structure.

Increasing salinity separated samples along the x-axis, with sites from the Murray Mouth on the far right and those from the South Lagoon on the far left of the dbRDA plot (Figure 24); which explained 27.1% of the total variation in the data. A further 14.5% was explained by changes in the percentage of fine sand and Chlorophyll-a in sediments, representing differences along the vertical axis of the plot

(Figure 24). Noonameena and the northern shore of Parnka Point contained macroinvertebrate assemblages more similar to the South Lagoon, while mudflats at Long Point and Mark Point were inhabited by similar macroinvertebrate assemblages as found in the Murray Mouth, which appeared to be affected mainly by salinity and sediment properties (Figure 24).





Figure 24. Distance-based redundancy analysis (dbRDA) plot with vector overlay showing environmental variables best explaining the pattern in macroinvertebrate community structure for the 2013/14 water release monitoring. The symbols represent the macroinvertebrate assemblage at each site for each sampling occasion (pooled across replicates). Symbols plotted close together represent similar macroinvertebrate communities. Symbols used here represent regions, whereby the North Lagoon was divided further into sub-regions due to the observed difference in macroinvertebrate communities occurring north and south of about Noonameena.


Assessing links between environmental variables and macroinvertebrates by region, the strongest explanations of macroinvertebrate communities by water or sediment properties were found for the South Lagoon and Murray Mouth (Figure 25). In the South Lagoon, the increase in salinity over the summer months accounted for the differentiation of macroinvertebrate assemblages at Villa de Yumpa in February and March 2014, whereas this site grouped closer to the southern shores of Parnka Point in December 2013. Differences over summer 2013/14 in the macroinvertebrate assemblage at Parnka Point South were more affected by grain size compositions (Figure 25). Overall, salinity gradients were the only significant contributor to differences in macroinvertebrate community structure in the South Lagoon (Pseudo- $F = 8.424$; $P_{(perm)} = 0.0304$), explaining nearly 68% of the total variation in the data.

For the Murray Mouth, the seasonal increase in salinity over summer accounted for differences in the macroinvertebrate community occurring in December 2013 from the one present in February and March 2014 (Figures 4 and 25), with salinity contributing significantly (38%, Pseudo- $F = 6.121$; $P_{(perm)} = 0.0005$) to the 40.9% of variability in macroinvertebrate communities explained by the

Murray Mouth



North Lagoon



South Lagoon




Figure 25. Distance-based redundancy analysis (dbRDA) plot with vector overlay showing environmental variables best explaining patterns in macroinvertebrate community structure for each region across the sampling occasions in 2013/14. Environmental variables at Parnka Point North (PaPN) were only determined in December 2013 during the TLM monitoring. Only two sites in the northern part of the South Lagoon were sampled.

measured environmental variables (Figure 25). Further differentiation of macroinvertebrate communities at the Murray Mouth sites over the 2013/14 survey period was mainly due to changes in microphytobenthic biomass (Chl-a, Pseudo- $F = 3.381$; $P_{(perm)} = 0.0062$) in the mudflat sediments, which were higher in February and March 2014 compared to December 2013, especially at Monument Road and Pelican Point (Appendix Figure A5). Salinity and Chl-a together explained about 58% of the variation in the macroinvertebrate community in the Murray Mouth over the 2013/14 survey.

In the North Lagoon, only 21.9% of the variation in macroinvertebrate communities was explained by environmental variables (Figure 25). This assessment is affected by the inclusion of Parnka Point North for December 2013. The sampling sites in the North Lagoon had very different salinities during sampling in December 2013 (Figure 4), and the hypersaline conditions at Parnka Point North and Noonameena at the time accounted for the different macroinvertebrate assemblage found. In February and March 2014, marine salinities occurred in the North Lagoon (Figure 4) and there was little time or site specific pattern for the macroinvertebrate assemblage (Figure 25). Neither salinity nor sediment grain size contributed significantly ($P_{(perm)} > 0.05$) to variation in macroinvertebrate communities in the North Lagoon.

Key Question 3) Are there similarities or differences in the community structure of macroinvertebrates across differing flow scenarios?

As described above (Key Question 1g), macroinvertebrate communities differed between the pre-drought, drought and flow periods (Figure 22), with indications for a return to pre-drought conditions in the Murray Mouth, or a possible change into a new state in the Coorong derived from trajectories (Appendix Figure A15). The macroinvertebrate communities found during the drought were highly dissimilar to those observed during the flow period in both the Murray Mouth and Coorong, mainly as additional species occurred as characteristic components of the post-flow community (Table 6).

For the Murray Mouth, the macroinvertebrate community in the flow period was characterised by a dominance of amphipods and chironomid larvae, yet in the recent survey further species characterised the community, such as the polychaetes *Simplisetia aequisetis* and *Boccardiella limnicola*. Amphipods and polychaetes, as well as the molluscs (*Salinator fragilis*, *Arthritica helmsi*), changed in abundance over the years and with it their contribution to defining communities. These species were part of a typical estuarine macroinvertebrate community as common in estuaries in southern Australia (Hirst 2004, Tweedley et al. 2012). The diversity of macroinvertebrates in the Murray Mouth is, however, much lower than in estuaries in West Australia or Victoria (Hirst 2004, Tweedley et al. 2012).

Multivariate canonical discriminant analysis of principle coordinates (CAP) highlighted the differences in communities between pre-drought, drought and flow conditions, with a clear and significant (Trace = 1.139; $P_{(perm)} = 0.001$) separation of communities from the three periods and the current monitoring. The squared canonical correlation of the first axis is high, indicating a good separation of samples (Figure 26). As already seen with the SIMPER analysis, flow years were characterised by the presence of chironomid larvae and amphipodes, while *Capitella* sp., *Simplisetia aequisetis* and *Arthritica helmsi* were more characteristic for the pre-drought and drought period (Figure 26).

Misclassification rate for this model was modest at 13%. Macroinvertebrate communities in the Murray Mouth during the 2013/14 monitoring were classified by the CAP model as belonging to a flow regime.

In the Coorong, chironomid larvae and *Capitella* sp. dominated the macroinvertebrate community in all periods, with amphipods being a further characteristic part of the community in the flow period and the recent monitoring (Table 6). As the community in the Coorong has been species poor, changes in abundances and rare findings of additional macroinvertebrates at single sampling occasions would account for the high dissimilarity across all time periods (Table 6). CAP analysis showed that in the Coorong, distinctions between macroinvertebrate communities during different flow regimes were less clear, but significant (Trace = 0.841; $P_{(perm)} = 0.007$). Communities in the drought and flow periods were on either end of the pre-drought period (Figure 26). The squared canonical correlation of the first axis was not as high as for the Murray Mouth region, but high enough to ensure a good representation of separation of communities based on flow regimes (Figure 26). Communities during flow years were characterised by the increased presence of amphipods and mysid shrimps (Figure 26). The misclassification rate for the Coorong model was 31.43%, higher than for the Murray Mouth. The macroinvertebrate community in the Coorong during the 2013/14 monitoring was classified by the CAP model as belonging to a flow regime.

Table 6. SIMPER results showing average similarities (light grey shading) and characteristic taxa for each flow regime (see Figure 21) as well as average dissimilarity (no shading) among communities for different flow regimes in a) the Murray Mouth, and b) Coorong Lagoons. Species are Cap: *Capitella* sp.; Sa: *Simplisetia aequisetis*; Sf: *Salinator fragilis*; Na: *Nephtys australiensis*; Bl: *Boccardiella limnicola*; Amp: Amphipoda; Ah: *Arthritica helmsi*; Chi: Chironomidae (larvae); Ost: Ostracoda; Eph: Ephidridae (larvae).

a) Murray Mouth	Pre-drought	Drought	Flow	2013-14	Characterising Taxa
Pre-drought	60.43				Ah; Sa; Cap; Amp
Drought	59.91	36.46			Cap; Sa; Sf; Na; Amp
Flow	67.71	75.16	42.63		Amp; Chi
2013-14	52.93	71.47	53.64	78.50	Amp; Sa; Chi; Bl

b) Coorong	Pre-drought	Drought	Flow	2013-14	Characterising Taxa
Pre-drought	21.97				Chi; Cap
Drought	73.16	37.78			Chi; Cap
Flow	76.03	78.10	27.55		Chi; Amp; Cap
2013-14	76.57	79.00	69.65	33.15	Chi; Cap; Eph; Ost; Amp




Figure 26. Canonical analysis of principle coordinates (CAP) plot showing macroinvertebrate communities when sampled in the a) Murray Mouth, and b) Coorong Lagoons during pre-drought, drought and flow years (see key in Figure 21 for monitoring events belonging to each flow regime).

5 Conclusions

In the fourth year since flows returned, the diversity and abundances of macroinvertebrates in the Murray Mouth and Coorong were showing stronger signs of recovery in late spring/early summer, followed by a drop in numbers of previously dominant species in the course of summer. Based on long-term comparisons of late spring/early summer surveys, a cycle of change was apparent for the Murray Mouth, where macroinvertebrate communities were becoming more similar to the pre-drought period, while a similar development in the Coorong was less pronounced. Whether the recovery will lead to a restoration of pre-drought conditions or to a new condition remains to be further investigated.

Recovery of estuarine ecosystems can take time (Duarte et al. 2014) and, as this monitoring showed, pre-drought condition have not yet been restored. A sequence of multiple stressors can affect the resilience of estuarine macroinvertebrates (Verdelhos et al. 2014) and the prolonged drought and intensive flow peak in 2010/11 may have affected the resilience of the Murray Mouth and Coorong. Freshwater flows can have effects similar to other disturbances and lead to a peak of opportunists in the early phases of recolonisation, followed by transitional periods to recovery (Pearson & Rosenberg 1978, Cañedo-Argüelles & Rieradevall 2010, Duggan et al. 2014). For future climate scenarios with increased frequencies of extreme events, findings from this monitoring will assist management considerations into the future.

Subtidal locations were only sampled on one occasion in the 2013/14 monitoring, and revealed higher abundances in sediments in shallow water. By not continuing subtidal sampling over summer, population size estimates or assessments of recruitment could not be given for some species as they were mostly or only recorded from subtidal sediments. For the records of presence and distribution of species such as some of the larger bivalves, the present monitoring has very likely underestimated the distribution ranges of macroinvertebrate species.

The decrease in abundances of amphipods and chironomid larvae in early 2014 exceeded smaller seasonal reductions in abundances seen in the previous surveys. Foraging depletion by shorebirds can reduce prey abundances in overwintering habitats (Saleem et al. 2014) and the effect of predation pressure from shorebirds in the Murray Mouth and Coorong merits investigation. Alternative explanations for the decrease could lie in the extreme heat wave in South Australia in January 2014, which could have exceeded the thermal tolerance of macroinvertebrate species (Poertner et al. 2002).

Polychaetes like *Simplisetia aequisetis* did not show such a pronounced reduction in abundances over summer, and their ability to dwell deeper in the sediment may have provided a thermal refuge. Their burrowing activity will further benefit the biogeochemical recovery of sediments in the Murray Mouth and Coorong, which have been anaerobic in many places and covered with filamentous macroalgae (Kristensen & Mikkelsen 2003, Lohrer et al. 2004). With more grazing snails and surface deposit-feeding bivalves present, the texture of the sediment-water/sediment-air interface will become roughened, which could further benefit exchange processes and improvements of sediment biogeochemistry (Maire et al. 2007). Such feedback mechanisms along the recovery pathway are of interest to understand the processes driving the resilience of the system.

6 Acknowledgements

This monitoring would not have been possible without further helping hands in the field and lab that were provided by Michael Drew, Shea Cameron, Sasha Whitmarsh and Stephanie Baggalley. Their assistance in collecting, sorting and identifying samples is greatly appreciated. For longer term comparisons, data obtained through The Living Murray Icon Site Monitoring, funded by the Murray-Darling Basin Authority, were used in this report. This project was funded by the Department of Environment, Water and Natural Resources and is part of the South Australian Government's *Murray Futures* program, funded in turn by the Australian Government's Water for the Future initiative. The entire CLLMM team are acknowledged for ongoing discussion and support of the project.

7 References

- Attrill, M.J., Rundle, S.D., 2002. Ecotone or Ecoline: Ecological Boundaries in Estuaries. *Estuarine, Coastal and Shelf Science* 55, 929-936.
- Baring, R., Dittmann, S., Dutton, A., Gannon, R., Cummings, S., Humphries, J., Hunt, T., 2009. Macrobenthic Survey 2008: Murray Mouth, Coorong and Lower Lakes Ramsar Site. Report for the South Australian Murray-Darling Basin Natural Resources Management Board, Adelaide, pp. 67 pp.
- Borja, A., Dauer, D.M., Elliott, M., Simenstad, C.A., 2010. Medium- and long-term recovery of estuarine and coastal ecosystems: patterns, rates and restoration effectiveness. *Estuaries and Coasts* 33, 1249-1260.
- Cañedo-Argüelles, M., Rieradevall, M., 2010. Disturbance caused by freshwater releases of different magnitude on the aquatic macroinvertebrate communities of two coastal lagoons. *Estuarine, Coastal and Shelf Science* 88(2), 190-198.
- Cummings, V.J., Thrush, S.F., 2004. Behavioural response of juvenile bivalves to terrestrial sediment deposits: implications for post-disturbance recolonisation. *Marine Ecology Progress Series* 278, 179-191.
- Dittmann, S., Nelson, M., 2007. Macrobenthic Survey 2006 in the Murray Mouth, Coorong and Lower Lakes Ramsar Site. Report for the Department for Environment and Heritage, Adelaide.
- Dittmann, S., Dutton, A., Earl, J., 2008. Macrobenthic Survey 2007 in the Murray Mouth, Coorong and Lower Lakes Ramsar Site. Report for the Department for Environment and Heritage, Adelaide.
- Dittmann, S., Cantin, A., Noble, W., Pocklington, J., 2006. Macrobenthic Survey 2004 in the Murray Mouth, Coorong and Lower Lakes Ramsar Site, with an Evaluation of Food Availability for Shorebirds and Possible Indicator Functions of Benthic Species., Adelaide, pp. 55 pp.
- Dittmann, S., Cantin, A., Imgraben, S., Ramsdale, T., 2006. Macrobenthic Survey 2005 in the Murray Mouth, Coorong and Lower Lakes Ramsar Site. Report for the Department for Environment and Heritage, Adelaide.
- Dittmann, S., Baggalley, S., Brown, E., Drew, M., Keuning, J., 2011. Benthic macroinvertebrate monitoring for the Goolwa Channel Water Level Management Project, year two, and Barrage Releases within the Coorong, Lower Lakes and Murray Mouth region. Report for the Department of Environment and Natural Resources. Flinders University, Adelaide.
- Dittmann, S., Ramsdale, T., Keuning, J., Navong, N., Baggalley, S., 2013. Benthic Macroinvertebrate Response Monitoring in the Coorong and Murray Mouth, 2012/13. Report for the Department of Environment, Water and Natural Resources, Adelaide.
- Dittmann, S., Baggalley, S., Baring, R., Brown, E., Gannon, R., Silvester, L., 2010. Macrobenthic survey 2009: Murray Mouth, Coorong and Lower Lakes Ramsar site. Report for the South Australian Murray Darling Basin Natural Resources Management Board, Adelaide.
- Dittmann, S., Ramsdale, T., Navong, N., Cummings, C., Baggalley, S., Keuning, J., 2013. Benthic macroinvertebrate survey 2012-13: Lower Lakes, Coorong and Murray Mouth Icon Site. , Report

- for the Department of Environment, Water and Natural Resources and the Murray-Darling Basin Authority. Flinders University of South Australia.
- Dittmann, S., Brown, E., Navong, N., Beyer, K., Silvester, L., Baggalley, S., Keuning, J., 2012. Macrobenthic invertebrate survey 2011-12: Lower Lakes, Coorong and Murray Mouth Icon Site. Report for the Department for Water and Murray-Darling Basin Authority. Flinders University of South Australia.
- Dolbeth, M., Cardoso, P.G., Ferreira, S.M., Verdelhos, T., Raffaelli, D., Pardal, M.A., 2007. Anthropogenic and natural disturbance effects on a macrobenthic estuarine community over a 10-year period. *Marine Pollution Bulletin* 54(5), 576-585.
- Duarte, C.M., Borja, A., Carstensen, J., Elliott, M., Krause-Jensen, D., Marbà, N., 2014. Paradigms in the Recovery of Estuarine and Coastal Ecosystems. *Estuaries and Coasts*, (in press).
- Duggan, M., Connolly, R.M., Whittle, M., Curwen, G., Burford, M.A., 2014. Effects of freshwater flow extremes on intertidal biota of a wet-dry tropical estuary. *Marine Ecology Progress Series* 502, 11-23.
- Elliot, M., Burdon, D., Hemingway, K.L., Apitz, S.E., 2007. Estuarine, coastal and marine ecosystem restoration: Confusing management and science - A revision of concepts. *Estuarine, Coastal and Shelf Science* 74, 349-366.
- Geddes, M.C., 1987. Changes in salinity and in the distribution of macrophytes, macrobenthos and fish in the Coorong lagoons, South Australia, following a period of River Murray flow. *Trans. R. Soc. S. Aus.* 111(4), 173-181.
- Geddes, M.C., Butler, A.J., 1984. Physicochemical and biological studies on the Coorong Lagoons, South Australia, and the effects of salinity on the distribution of the macrobenthos. *Transactions of the Royal Society of South Australia* 108, 51-62.
- Gillet, P., Torresani, S., 2003. Structure of the population and secondary production of *Hediste diversicolor* (O.F. Müller, 1776), (Polychaeta, Nereidae) in the Loire estuary, Atlantic Coast, France. *Estuarine, Coastal and Shelf Science* 56(3-4), 621-628.
- Günther, C.-P., 1992. Dispersal of intertidal invertebrates: A strategy to react to disturbances of different scales? *Netherlands Journal of Sea Research* 30, 45-56.
- Hirst, A., 2004. Broad-scale environmental gradients among estuarine benthic macrofaunal assemblages of south-eastern Australia: implications for monitoring estuaries. *Marine and Freshwater Research* 55, 79-92.
- Hummel, H., Meijeboom, A., De Wolf, L., 1986. The effects of extended periods of drainage and submersion on condition and mortality of benthic animals. *Hydrobiologia* 103, 251-266.
- Hummel, H., Fortuin, A.W., De Wolf, L., Meijeboom, A., 1988. Mortality of intertidal benthic animals after a period of prolonged emersion. *Journal of Experimental Marine Biology and Ecology* 121, 247-254.
- Kangas, M.I., Geddes, M.C., 1984. The effects of salinity on the distribution of amphipods in the Coorong, South Australia, in relation to their salinity tolerance. *Transactions of the Royal Society of South Australia* 108(3), 139-145.
- Keuning, J., Brown, E., Navong, N., Beyer, K., Baggalley, S., Dittmann, S., 2012. Benthic macroinvertebrate response monitoring in the Murray Mouth and Coorong, 2011/12. Report for the Department for Environment and Natural Resources, Adelaide.
- Kristensen, E., Mikkelsen, O.L., 2003. Impact of the burrow-dwelling polychaete *Nereis diversicolor* on the degradation of fresh and aged macroalgal detritus in a coastal marine sediment. *Marine Ecology Progress Series* 265, 141-153.
- Leblanc, M., Tweed, S., van Dijk, A., Timbal, B., 2012. A review of historic and future hydrological changes in the Murray-Darling Basin. *Global and Planetary Change* 80-81, 226-246.
- Levin, L.A., Boesch, D.F., Covich, A., Dahm, C., Erséus, C., Ewel, K.C., Kneib, R.T., Moldenke, A., Palmer, M.A., Snelgrove, P.V.R., Strayer, D., Weslawski, J.M., 2001. The function of marine critical transition zones and the importance of sediment biodiversity. *Ecosystems* 4, 430-451.
- Lohrer, A.M., Thrush, S.F., Gibbs, M.M., 2004. Bioturbators enhance ecosystem function through complex biogeochemical interactions. *Nature* 431, 1092-1095.

- Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H., Cooke, R.G., Kay, M.C., Kidwell, S.M., Kirby, M.X., Peterson, C.H., Jackson, J.B.C., 2006. Depletion, Degradation, and Recovery Potential of Estuaries and Coastal Seas. *Science* 312(5781), 1806-1809.
- Maire, O., Duchêne, J.C., Grémare, A., Malyuga, V.S., Meysman, F.J.R., 2007. A comparison of sediment reworking rates by the surface deposit-feeding bivalve *Abra ovata* during summertime and wintertime, with a comparison between two models of sediment reworking. *Journal of Experimental Marine Biology and Ecology* 343(1), 21-36.
- Marques, L., Carrico, A., Bessa, F., Gaspar, R., Neto, J.M., Patricio, J., 2013. Response of intertidal macrobenthic communities and primary producers to mitigation measures in a temperate estuary. *Ecological Indicators* 25, 10-22.
- Oishi, K., Saigusa, M., 1999. Rhythmic patterns of abundance in small sublittoral crustaceans: variety in the synchrony with day/night and tidal cycles. *Marine Biology* 133(2), 237-247.
- Olafsson, E.B., Peterson, C.H., Ambrose, W.G.j., 1994. Does recruitment limitation structure populations and communities of macro-invertebrates in marine soft sediments: the relative significance of pre- and post-settlement processes. *Oceanography and Marine Biology: an Annual Review* 32, 65-109.
- Pearson, T.H., Rosenberg, R., 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment *Oceanography Marine Biology Annual Review* 16, 229-311.
- Poertner, H.O., 2002. Climate variations and the physiological basis of temperature depend on biogeography: systemic to molecular hierarchy of thermal tolerance in animals *Comparative Biochemistry and Physiology Part A* 132, 739-761.
- Potter, I.C., Chuwen, B.M., Hoeksema, S.D., Elliott, M., 2010. The concept of an estuary: A definition that incorporates systems which can become closed to the ocean and hypersaline. *Estuarine, Coastal and Shelf Science* 87, 497-500.
- Ramskov, T., Forbes, V.E., 2008. Life history and population dynamics of the opportunistic polychaete *Capitella* sp. I in relation to sediment organic matter. *Marine Ecology Progress Series* 369, 181-1192.
- Salem, M.V.A., van der Geest, M., Piersma, T., Saoud, Y., van Gils, J.A., 2014. Seasonal changes in mollusc abundance in a tropical intertidal ecosystem, Banc d'Arguin (Mauritania): Testing the 'depletion by shorebirds' hypothesis. *Estuarine, Coastal and Shelf Science* 136, 26-34.
- Sebasvari, Z., Esser, F., Harder, T., 2006. Sediment-associated cues for larval settlement of the infaunal spionid polychaetes *Polydora cornuta* and *Streblospio benedicti*. *Journal of Experimental Marine Biology and Ecology* 337, 109-120.
- Thrush, S.F., Whittatch, R.B., Pridmore, R.D., Hewitt, J.E., Cummings, V.J., Wilkinson, M.R., 1996. Scale-dependent recolonization: The role of sediment stability in a dynamic sandflat habitat. *Ecology* 77(8), 2472-2487.
- Tsutsumi, H., 1990. Population persistence of *Capitella* sp. (Polychaeta; Capitellidae) on a mud flat subject to environmental disturbance by organic enrichment. *Marine Ecology Progress Series* 63, 147-156.
- Tweedley, J.R., Warwick, R.M., Valesini, F.J., Platell, M.E., Potter, I.C., 2012. The use of benthic macroinvertebrates to establish a benchmark for evaluating the environmental quality of microtidal, temperate southern hemisphere estuaries. *Marine Pollution Bulletin* 64(6), 1210-1221.
- Verdelhos, T., Cardoso, P.G., Dolbeth, M., Pardal, M.A., 2014. Recovery trends of *Scrobicularia plana* populations after restoration measures, affected by extreme climate events. *Marine Environmental Research* 98, 39-48.
- Verdonschot, P.F.M., Spears, B.M., Feld, C.K., Brucet, S., Keizer-Vlek, H., Borja, A., Elliott, M., Kernan, M., Johnson, R.K., 2013. A comparative review of recovery processes in rivers, lakes, estuarine and coastal waters. *Hydrobiologia* 704(1), 453-474.
- Verissimo, H., Bremner, J., Garcia, C., Patricio, J., van der Linden, P., Marques, J.C., 2012. Assessment of the subtidal macrobenthic community functioning of a temperate estuary following environmental restoration. *Ecological Indicators* 23, 312-322.
- Whitfield, A.K., Elliott, M., Basset, A., Blaber, S.J.M., West, R.J., 2012. Paradigms in estuarine ecology - A review of the Remane diagram with a suggested revised model for estuaries. *Estuarine, Coastal and Shelf Science* 97, 78-90.

Whitlatch, R.B., Lohrer, A.M., Thrush, S.F., Pridmore, R.D., Hewitt, J.E., Cummings, V.J., Zajac, R.N., 1998. Scale-dependent benthic recolonization dynamics: life stage-based dispersal and demographic consequences. *Hydrobiologia* 375/376, 217-226.