



Other pest plant control options sheets available:

- African boxthorn
- African Lovegrass
- African Rue
- Alkali Sida
- Alligator Weed
- Athel pine
- Bathurst Burr
- Blackberry
- Bladder Campion
- Boneseed
- Bridal Creeper
- Calomba Daisy
- Caltrop
- Cape tulip
- Coolatai Grass
- Creeping Knapweed
- Cutleaf Mignonette
- Dog Rose
- Gorse
- Hoary Cress
- Horehound
- Innocent Weed
- Khaki Weed
- Lincoln Weed
- Noogoora Burr
- Pepper Trees
- Perennial Ragweed
- Prickly pear/ Wheel cactus
- Salvinia
- Silverleaf Nightshade
- Water Hyacinth
- Water Primrose
- Wild artichoke
- Willows
- Yellow Burrweed

For further info please contact:

Ph. (08) 8532 1432



Hoary Cress

Cardaria draba

AIM: To destroy plants and prevent any further seed set

Control Options	Chemical	Rate	Comments
Spot Spray	Tordon [®] 75D	1.3L/100L	Soil residual. Do not use near desirable vegetation or in waterways. Do not apply near susceptible crops. May require repeat applications.
	Weedmaster [®] Duo	500ml/100L	Apply from late July to September when plants are in rosette to flowering stage
Boom Spray	Weedmaster [®] Duo	1.5L/ha	Apply from late July to September when plants are in rosette to flowering stage
	Broadside [™]	1L/ha	Apply when at 2 to 4 leaf stage.

Always follow safe use instructions on herbicide labels. Refer to product label for full conditions of use and application instructions.

These options have been compiled from herbicide manufacturer labels or previous experience and the South Australian Murray-Darling Basin Natural Resources Management Board (SA MDB NRM Board) holds no responsibility for their effectiveness. Permits may be required for any off label use.

For specific advice on the use of these herbicides and options please contact the SA MDB NRM Board Officers. In certain circumstances these options may not be suitable.