

SAND OVER ACIDIC CLAY

General Description: *Medium to thick sandy surface soil with moderate to heavy ironstone gravel overlying a brown, yellow and red clay grading to highly weathered kaolinised soft sandstone.*

Landform: Upper slopes and crests of undulating rises and low hills associated with ancient glacial valleys

Substrate: Soft red, grey and yellow mottled kaolinised sandstone

Vegetation: Eucalyptus baxteri / Euc. fasciculosa scrub



Type Site:	Site No.:	CH015	1:50,000 mapsheet:	6627-3 (Willunga)
	Hundred:	Nangkita	Easting:	288700
	Section:	232	Northing:	6081200
	Sampling date:	29/07/92	Annual rainfall:	790 mm average

Upper slope of undulating low hills, 9% slope. Soft surface with a trace of ironstone.

Soil Description:

<i>Depth (cm)</i>	<i>Description</i>
0-12	Very dark grey soft light sandy loam. Clear to:
12-31	Very pale brown soft loamy sand. Abrupt to:
31-39	Reddish yellow soft loamy sand with up to 50% ironstone and sandstone. Abrupt to:
39-54	Brown friable light medium clay. Clear to:
54-75	Brownish yellow and light brown friable light clay. Gradual to:
75-150	Brown, red and yellow firm massive fine sandy clay. Diffuse to:
150-200	Yellow, white and red massive soft sandstone.



Classification: Bleached, Eutrophic, Brown Kurosol; thick, non-gravelly, loamy / clayey, very deep



Summary of Properties

- Drainage:** Well drained. The soil is unlikely to remain wet for more than a few days.
- Fertility:** The natural fertility of the soil is low to moderate, as indicated by the exchangeable cation data, although the surface relies on its organic matter content for satisfactory nutrient retention. The data indicate that although cation ratios are good, there are marginal deficiencies of calcium, magnesium and potassium. Manganese and copper are possibly deficient as well.
- pH:** Acidic at the surface, strongly acidic at base. Dolomitic lime is needed for correction.
- Rooting depth:** 150 cm at type site, although density is low from 75 cm.
- Barriers to root growth:**
- Physical:** There are no physical barriers to root growth.
 - Chemical:** Low pH, with possible associated aluminium toxicity, and nutrient leaching are the main limitations. Ironstone gravel tends to fix phosphorus, indicating higher rates are required.
- Waterholding capacity:** 180 mm in rootzone, but only about two thirds of this is effectively available because of poor root distribution.
- Seedling emergence:** Good.
- Workability:** Good, except for abrasive effects of ironstone.
- Erosion Potential**
- Water:** Moderate because of the slope and the low permeability of the clayey subsoil.
 - Wind:** Moderately low. The light surface will erode if exposed.

Laboratory Data

Depth cm	pH H ₂ O	pH CaCl ₂	CO ₃ %	EC1:5 dS/m	ECe dS/m	Org.C %	Avail. P mg/kg	Avail. K mg/kg	SO ₄ mg/kg	Boron mg/kg	Trace Elements mg/kg (DTPA)				CEC cmol (+)/kg	Exchangeable Cations cmol(+)/kg				ESP
											Cu	Fe	Mn	Zn		Ca	Mg	Na	K	
Paddock	5.9	5.5	-	0.08	0.37	3.9	70	190	-	1.0	0.9	99	2.3	2.6	7.4	4.9	1.4	<0.1	0.36	<1.5
0-12	6.0	5.6	-	0.07	0.26	3.4	12	120	-	0.7	0.2	103	1.4	1.7	5.7	4.0	1.4	<0.1	0.21	<2
12-31	5.6	4.8	-	0.04	0.14	0.4	5	120	-	0.5	<0.1	94	<0.1	<0.1	1.7	0.6	0.2	<0.1	0.25	na
31-39	5.3	4.7	-	0.06	0.16	0.8	8	220	-	0.8	0.8	254	0.2	0.5	4.4	1.4	0.5	0.12	0.50	2.7
39-54	5.3	4.6	-	0.06	0.14	1.1	<2	360	-	2.0	0.1	53	0.1	<0.1	11.1	3.3	2.9	0.29	0.94	2.6
54-75	5.5	5.2	-	0.06	0.16	0.4	<2	110	-	1.6	<0.1	9	<0.1	<0.1	8.9	2.3	4.0	0.25	0.28	2.8
75-150	5.2	5.0	-	0.05	0.27	<0.1	<2	25	-	0.8	<0.1	2	<0.1	<0.1	2.5	0.6	1.6	0.12	0.05	na
150-200	4.9	4.4	-	0.04	0.15	<0.1	<2	17	-	0.6	<0.1	2	<0.1	<0.1	2.2	0.4	1.1	0.12	<0.05	na

Note: Paddock sample bulked from 20 cores (0-10 cm) taken around the pit.

* EDTA trace element analyses for "paddock" sample.

CEC (cation exchange capacity) is a measure of the soil's capacity to store and release major nutrient elements.

ESP (exchangeable sodium percentage) is derived by dividing the exchangeable sodium value by the CEC.

Further information: [DEWNR Soil and Land Program](#)

